



UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS  
International General Certificate of Secondary Education

---

**ADDITIONAL MATHEMATICS (US)**

**0459/01**

Paper 1

**For Examination from 2013**

SPECIMEN MARK SCHEME

**2 hours**

---

**MAXIMUM MARK: 80**

---

This document consists of **9** printed pages and **1** blank page.

## Mark Scheme Notes

- Marks are of the following three types:

**M** Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark, and in some cases an M mark can be implied from a correct answer.

**A** Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

**B** Accuracy mark for a correct result or statement independent of method marks.

- When a part of a question has two or more “method” steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB is used to indicate that a particular M or B mark is dependent on an earlier M or B mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.

- It implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously “correct” answers or results obtained from incorrect working.

- Note. B2 or A2 means that the candidate can earn 2 or 0.  
B2, 1, 0 means that the candidate can earn anything from 0 to 2. –1 each error. A mark is deducted from the total mark available up to the maximum mark available for that question. The minimum mark awarded is zero e.g., if a candidate makes 3 errors in a question worth 2 marks they score zero.

- The following abbreviations may be used in a mark scheme.

**AG** ‘Answer given’ on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid).

**cao** ‘Correct answer only’ (emphasizing that no “follow through” from a previous error is allowed).

**isw** ‘Ignore subsequent working’.

**oe** ‘Or equivalent’.

**sc** ‘Special case’. Awarded for some questions where e.g., the candidate has not used the method specified but a different, correct, method leading to the correct answer.

**soi** ‘Seen or implied’.

| Question | Answer  | Mark   | Guidance   |
|----------|---|--|--|
| 1        | $(x + 2)^2 + (y - 8)^2 = 7^2$<br>or $x^2 + y^2 + 4x - 16y + 19 = 0$   | <b>B2</b><br><br>[2]                                     | <b>B1</b> for $(x + 2)^2 + (y - 8)^2 = 7^2$<br>or $x^2 + 4x + y^2 + 16y = 49$ oe<br><b>B1</b> for $x^2 + y^2 + 4x - 16y + 19 = 0$ oe           |
| 2 (a)    | Any two valid reasons<br>e.g.<br>Size of population may make selection of every item impossible<br>Gathering information may necessitate destruction of items e.g.<br>life of a battery             | <b>B1 + B1</b>   |  |
| (b)      | Every member of population has the same chance of being selected at every stage if random and choice of 1 <sup>st</sup> item immediately rules out approximately 90% of the remaining population oe | <b>B2, 1, 0</b><br><br>[4]                               | Explanation must incorporate the essential idea of random sampling.  |
| 3        | $(z_1 =) \frac{3 + \sqrt{7i^2}}{2}$<br><br>$(z_1 =) \frac{3 + i\sqrt{7}}{2}$<br><br>$(z_2 =) \frac{3 - i\sqrt{7}}{2}$   | <b>M1</b><br><br><b>A1</b><br><br><b>B1ft</b><br><br>[3] | allow $z =$ ; allow $\frac{3 \pm \sqrt{7i^2}}{2}$<br><br>ft the complex conjugate of their $z_1$<br>$\frac{3 \pm i\sqrt{7}}{2}$ scores 3 marks |

|   |   |  |   |
|---|---|--|---|
| 4 | $(PS)^2 = (x-6)^2 + (y-1)^2$ $(x-6)^2 + (y-1)^2 = (x+1)^2$ $x^2 - 12x + 36 + y^2 - 2y + 1 = x^2 + 2x + 1$ $y(y-2) = 14x - 36 \quad \text{AG}$   | <b>B1</b><br><b>M1</b><br><br><b>M1</b><br><br><b>A1</b><br><b>[4]</b>                               |   |
| 5 | $(5 + 2\sqrt{3})^2 = 37 + 20\sqrt{3}$ $\frac{(37 + 20\sqrt{3})}{2 + \sqrt{3}} \times \frac{2 - \sqrt{3}}{2 - \sqrt{3}}$ $14 + 3\sqrt{3}$  | <b>B1</b><br><br><b>M1</b><br><br><br><b>A1+A1</b><br><b>[4]</b>                                     | <p>Seen anywhere</p> <p>Or <b>B1</b> for a correct pair of simultaneous equations <math>37 = 2p + 3q</math> and <math>20 = p + 2q</math></p> <p>and <b>M1</b> for attempting to solve their equations either by elimination or substitution, condone one error.</p> <p>Answer only scores zero.</p>   |
| 6 | <p>Proving triangle <i>AED</i> congruent to triangle <i>CFB</i><br/> <math>AD = BC</math> (parallelogram)<br/> <math>ED = FB</math> (given)<br/> <math>\angle ADE = \angle CBF</math> (alternate angles are equal)<br/> <math>\triangle AED \equiv \triangle CFB</math> (SAS)</p> <p><math>\angle AED = \angle CFB</math> (corresponding angles of congruent triangles)<br/> <math>\angle AEF = \angle CFE</math> (each equal to <math>180 - \angle AED</math>)</p> <p>Thus alternate angles are equal</p> <p><math>AE = FC</math> (corresponding sides of congruent triangles)</p> | <b>B3, 2, 1, 0</b><br><br><br><br><br><br><br><br><br><b>DB1</b><br><br><b>DB1</b><br><br><b>[5]</b> | <p>Or triangle <i>DEC</i> congruent to <i>BFA</i><br/> <math>AB = DC</math> (parallelogram)</p> <p><math>\angle ABF = \angle CDE</math> (alternate angles are equal)<br/> <math>\triangle ABF \equiv \triangle CDE</math> (SAS)<br/>         Must have reasons</p> <p><math>\angle AFB = \angle ECD</math> (corresponding angles of congruent triangles)<br/> <math>\angle AFE = \angle FEC</math> (each equal to <math>180 - \angle AFB</math>)</p> <p>Thus alternate angles are equal</p> <p><math>AF = EC</math> (corresponding sides of congruent triangles)<br/>         Other valid proofs should be awarded appropriate credit</p> |

|   |   |   |   |
|---|---|---|---|
| 7 | $\mathbf{A}^{-1} = k \begin{pmatrix} 1 & 3 \\ -1 & 2 \end{pmatrix}$ $k = \frac{1}{5}$ $\mathbf{A}^2 = \begin{pmatrix} 2 & -3 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 2 & -3 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & -9 \\ 3 & -2 \end{pmatrix}$ $\mathbf{B} = 2 \times \text{their} \begin{pmatrix} 1 & 3 \\ -1 & 2 \end{pmatrix} - \text{their} \begin{pmatrix} 1 & -9 \\ 3 & -2 \end{pmatrix}$ $\begin{pmatrix} 1 & 15 \\ -5 & 6 \end{pmatrix}$ | <p><b>B1</b></p> <p><b>B1</b></p> <p><b>M1</b></p> <p><b>A1</b></p><br><p><b>M1</b></p><br><p><b>A1</b></p> <p><b>[6]</b></p> | <p>attempt to multiply with at least two elements correct<br/>correct</p> |
| 8 | <p><b>(i)</b> <math>0.97 \times 0.04</math><br/><math>0.05 \times 0.96</math><br/>Summing their products<br/><math>0.0868</math></p> <p><b>(ii)</b> their <math>\frac{0.0388}{0.0868}</math><br/><math>0.447(00\dots)</math> A.G.</p>   | <p><b>M1</b></p> <p><b>M1</b></p> <p><b>M1</b></p> <p><b>A1</b></p><br><p><b>M1</b></p> <p><b>A1</b></p> <p><b>[6]</b></p>    |   |

|    |   |   |   |
|----|---|---|---|
| 9  | Eliminate $x$ or $y$<br>$4x^2 + 4x - 15 = 0$ or $4y^2 - 28y + 33 = 0$<br>Factorise 3 term quadratic<br>$x = \frac{3}{2}$ and $-\frac{5}{2}$<br>$y = \frac{11}{2}$ and $\frac{3}{2}$<br>$\sqrt{4^2 + 4^2}$<br>$\sqrt{32}$ or $4\sqrt{2}$ or 5.66 | <b>M1</b><br><b>A1</b><br><b>M1</b><br><br><b>A1</b><br><br><b>A1</b><br><br><b>M1</b><br><b>A1</b><br><br><b>[7]</b> |   |
| 10 | (i) $m_{AB} = \frac{1}{5}$<br>Uses $m_1 m_2 = -1$ ( $= m_{BC} = -5$ )<br>$BC: y - 5 = -5(x - 6)$ or $5x + y = 35$<br><br>$C(7,0)$<br>$CD: y - 0 = \frac{1}{5}(x - 7)$ oe<br><br>(ii) $D(1, -1.2)$   | <b>B1</b><br><br><b>M1</b><br><b>M1</b><br><br><b>A1</b><br><b>A1ft</b><br><br><b>B1ft</b><br><br><b>[6]</b>          | or gradient $BC = \frac{5}{6 - x_c} = -5$<br><br>ft their $C$ and $m_{AB}$<br><br>ft their equation of $CD$ |

|  |   |   |  |
|--|---|---|--|
| <p><b>11 (a)</b></p> <p><b>(b)</b></p> | <p><math>\sin x = 2\cos x</math><br/> <math>\tan x = 2</math><br/> 63.4<br/> 243.4</p> <p><math>2(1 - \cos^2 y) + 3\cos y = 0</math><br/> <math>2\cos^2 y - 3\cos y - 2 = 0</math><br/> <math>(2\cos y + 1)(\cos y - 2) = 0</math><br/> <math>\cos y = -\frac{1}{2}</math><br/> 120<br/> 240</p>  | <p><b>M1</b><br/> <b>M1</b><br/> <b>A1</b><br/> <b>A1</b></p> <p><b>M1</b></p> <p><b>M1</b></p> <p><b>A1</b><br/> <b>A1</b></p> <p><b>[8]</b></p> | <p>or correct use of quadratic formula or completing the square<br/> extra solutions within range <math>-1</math> (once each part)</p> |
| <p><b>12</b></p>                       | <p><math>(1200\mathbf{i} + 240\mathbf{j}) \div 4</math><br/> their <math>(300\mathbf{i} + 60\mathbf{j}) - (260\mathbf{i} + 156\mathbf{j})</math><br/> <math>40\mathbf{i} - 96\mathbf{j}</math></p> <p><math>\sqrt{40^2 + 96^2}</math><br/> 104</p> <p><math>\tan^{-1}\left(\frac{96}{40}\right)</math> or <math>\tan^{-1}\left(\frac{96}{40}\right)</math><br/> 157(.4)</p> | <p><b>M1</b><br/> <b>M1</b><br/> <b>A1</b></p> <p><b>M1</b><br/> <b>A1</b></p> <p><b>M1</b></p> <p><b>A1</b></p> <p><b>[7]</b></p>                | <p>clear indication of direction</p>   |

| <b>13</b>   | <b>(i)</b>  | $4\pi, 16\pi, 36\pi$<br>$4\pi, 16\pi - 4\pi, 36\pi - 16\pi$<br>$4\pi, 12\pi, 20\pi$  | <b>B1</b>                           |                |    |   |    |            |               |                |                |                |                 |  |
|-------------|---|--|-------------------------------------|----------------|----|---|----|------------|---------------|----------------|----------------|----------------|-----------------|--|
|             | <b>(ii)</b>   | $\frac{1}{9}$ soi<br>$\frac{1}{12}$  | <b>M1</b><br><b>B1</b><br><b>B1</b> |                |    |   |    |            |               |                |                |                |                 |  |
|             | <b>(iii)</b>  | <table border="1"> <thead> <tr> <th><math>S</math></th> <th>0</th> <th>3</th> <th>6</th> <th>12</th> </tr> </thead> <tbody> <tr> <td><math>P(S = s)</math></td> <td><math>\frac{1}{4}</math></td> <td><math>\frac{5}{12}</math></td> <td><math>\frac{3}{12}</math></td> <td><math>\frac{1}{12}</math></td> </tr> </tbody> </table> | $S$                                 | 0              | 3  | 6 | 12 | $P(S = s)$ | $\frac{1}{4}$ | $\frac{5}{12}$ | $\frac{3}{12}$ | $\frac{1}{12}$ | <b>B2, 1, 0</b> |  |
|             | $S$   | 0  | 3                                   | 6              | 12 |   |    |            |               |                |                |                |                 |  |
| $P(S = s)$  | $\frac{1}{4}$   | $\frac{5}{12}$   | $\frac{3}{12}$                      | $\frac{1}{12}$ |    |   |    |            |               |                |                |                |                 |  |
| <b>(iv)</b> | their $0 \times \frac{1}{4} + 3 \times \frac{5}{12} + 6 \times \frac{3}{12} + 12 \times \frac{1}{12}$ soi<br><br>3.75<br>75 | <b>M1</b><br><br><b>A1</b><br><b>A1 ft</b><br><b>[9]</b>   | ft their $E(S)$                     |                |    |   |    |            |               |                |                |                |                 |  |



|  |  |   |   |
|--|--|---|---|
| <p><b>14 (a) (i)</b></p> <p><b>(ii)</b></p> <p><b>(b) (i)</b></p> <p><b>(ii)</b></p> | $fg(x) = 3 - \frac{x}{x+2}$ $3 - \frac{x}{x+2} = 10$ <p><math>3(x+2) - x = 10(x+2)</math> or better<br/>leading to <math>x = -1.75</math></p> $h(x) > 4$ $h^{-1}(x) = e^{x-4}$ $h^{-1}(9) = e^5 (\approx 148)$ | <p><b>B1</b></p> <p><b>M1</b><br/><b>A1</b></p> <p><b>B1</b></p> <p><b>M1</b><br/><b>A1</b></p> | <p>for dealing with fraction appropriately state this mathematically</p> <p>for attempting to obtain inverse function</p> <p>or <b>M1</b> for <math>4 + \ln x = 9</math> and<br/><b>A1</b> for <math>x = e^5 (\approx 148)</math></p> |
| <p><b>(iii)</b></p>  | <p>correct graphs<br/>idea of symmetry</p>   | <p><b>B1 + B1</b><br/><b>B1</b><br/>[9]</p>   | <p>B1 for each curve</p>  |
|  |  | <p>[80]</p>   |   |

