

**Mathematics (MEI)**

Advanced GCE 4753

Methods for Advanced Mathematics (C3)

**Mark Scheme for June 2010**

---

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of pupils of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, OCR Nationals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by Examiners. It does not indicate the details of the discussions which took place at an Examiners' meeting before marking commenced.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the Report on the Examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.


© OCR 2010

Any enquiries about publications should be addressed to:

OCR Publications  
PO Box 5050  
Annesley  
NOTTINGHAM  
NG15 0DL

Telephone: 0870 770 6622  
Facsimile: 01223 552610  
E-mail: [publications@ocr.org.uk](mailto:publications@ocr.org.uk)

## Section A

<p><b>1</b></p> $\int_0^{\pi/6} \cos 3x dx = \left[ \frac{1}{3} \sin 3x \right]_0^{\pi/6}$ $= \frac{1}{3} \sin \frac{\pi}{2} - 0$ $= 1/3$	<p>M1 B1 A1cao [3]</p>	<p><math>k \sin 3x, k &gt; 0, k \neq 3</math> <math>k = (\pm)1/3</math> 0.33 or better</p>	<p>or M1 for <math>u = 3x \Rightarrow \int \frac{1}{3} \cos u du</math> condone <math>90^\circ</math> in limit or M1 for <math>\left[ \frac{1}{3} \sin u \right]</math> so: <math>\sin 3x</math>: M1B0, <math>-\sin 3x</math>: M0B0, <math>\pm 3 \sin 3x</math>: M0B0, <math>-1/3 \sin 3x</math>: M0B1</p>
<p><b>2</b></p> <p><math>f(x) =  x+1 </math>      <math>g(x) =  x +1</math></p> 	<p>B1 B1 B1 B1 [4]</p>	<p>soi from correctly-shaped graphs (i.e. without intercepts) graph of <math> x+1 </math> only graph of <math> x +1</math></p>	<p>but must indicate which is which bod gf if negative <math>x</math> values are missing 'V' shape with <math>(-1, 0)</math> and <math>(0, 1)</math> labelled 'V' shape with <math>(0, 1)</math> labelled <math>(0, 1)</math></p>
<p><b>3(i)</b></p> $y = (1+3x^2)^{1/2}$ $\Rightarrow \frac{dy}{dx} = \frac{1}{2}(1+3x^2)^{-1/2} \cdot 6x$ $= 3x(1+3x^2)^{-1/2}$	<p>M1 B1 A1 [3]</p>	<p>chain rule <math>\frac{1}{2} u^{-1/2}</math> o.e., but must be '3'</p>	<p>can isw here</p>
<p><b>(ii)</b></p> $y = x(1+3x^2)^{1/2}$ $\Rightarrow \frac{dy}{dx} = x \cdot \frac{3x}{\sqrt{1+3x^2}} + 1 \cdot (1+3x^2)^{1/2}$ $= \frac{3x^2 + 1 + 3x^2}{\sqrt{1+3x^2}}$ $= \frac{1+6x^2}{\sqrt{1+3x^2}} *$	<p>M1 A1ft M1 E1 [4]</p>	<p>product rule ft their <math>dy/dx</math> from (i) common denominator or factoring <math>(1+3x^2)^{-1/2}</math> www</p>	<p>must show this step for M1 E1</p>

<p><b>4</b> <math>p = 100/x = 100x^{-1}</math>  <math>\Rightarrow dp/dx = -100x^{-2} = -100/x^2</math></p> <p><math>dp/dt = dp/dx \times dx/dt</math>  <math>dx/dt = 10</math></p> <p>When <math>x = 50</math>, <math>dp/dx = (-100/50^2)</math>  <math>\Rightarrow dp/dt = 10 \times -0.04 = -0.4</math></p>	<p>M1 A1 M1 B1 M1dep A1cao [6]</p>	<p>attempt to differentiate  <math>-100x^{-2}</math> o.e.  o.e. soi  soi  substituting <math>x = 50</math> into their <math>dp/dx</math> dep 2<sup>nd</sup> M1  o.e. e.g. decreasing at 0.4</p>	<p>condone poor notation if chain rule correct  or <math>x = 50 + 10t</math> B1  <math>\Rightarrow P = 100/x = 100/(50 + 10t)</math>  <math>\Rightarrow dP/dt = -100(50 + 10t)^{-2} \times 10 = -1000/(50 + 10t)^2</math> M1  A1  When <math>t = 0</math>, <math>dP/dt = -1000/50^2 = -0.4</math> A1</p>
<p><b>5</b> <math>y^3 = xy - x^2</math>  <math>\Rightarrow 3y^2 dy/dx = x dy/dx + y - 2x</math></p> <p><math>\Rightarrow 3y^2 dy/dx - x dy/dx = y - 2x</math>  <math>\Rightarrow (3y^2 - x) dy/dx = y - 2x</math>  <math>\Rightarrow dy/dx = (y - 2x)/(3y^2 - x) *</math></p> <p>TP when <math>dy/dx = 0 \Rightarrow y - 2x = 0</math>  <math>\Rightarrow y = 2x</math>  <math>\Rightarrow (2x)^3 = x \cdot 2x - x^2</math>  <math>\Rightarrow 8x^3 = x^2</math>  <math>\Rightarrow x = 1/8</math> *(or 0)</p>	<p>B1 B1  M1 E1  M1 M1  E1 [7]</p>	<p><math>3y^2 dy/dx</math>  <math>x dy/dx + y - 2x</math></p> <p>collecting terms in <math>dy/dx</math> only</p> <p>or <math>x = 1/8</math> and <math>dy/dx = 0 \Rightarrow y = 1/4</math>  or <math>(1/4)^3 = (1/8)(1/4) - (1/8)^2</math>  or verifying e.g. <math>1/64 = 1/64</math></p>	<p>must show '<math>x dy/dx + y</math>' on one side</p> <p>or <math>x = 1/8 \Rightarrow y^3 = (1/8)y - 1/64</math> M1  verifying that <math>y = 1/4</math> is a solution (must show evidence*) M1  <math>\Rightarrow dy/dx = (1/4 - 2(1/8))/(...) = 0</math> E1  *just stating that <math>y = 1/4</math> is M1 M0 E0</p>
<p><b>6</b> <math>f(x) = 1 + 2 \sin 3x = y \quad x \leftrightarrow y</math>  <math>x = 1 + 2 \sin 3y</math>  <math>\Rightarrow \sin 3y = (x - 1)/2</math>  <math>\Rightarrow 3y = \arcsin [(x - 1)/2]</math>  <math>\Rightarrow y = \frac{1}{3} \arcsin \left[ \frac{x-1}{2} \right]</math> so <math>f^{-1}(x) = \frac{1}{3} \arcsin \left[ \frac{x-1}{2} \right]</math></p> <p>Range of <math>f</math> is <math>-1</math> to <math>3</math>  <math>\Rightarrow -1 \leq x \leq 3</math></p>	<p>M1 A1 A1  A1  M1 A1 [6]</p>	<p>attempt to invert</p> <p>must be <math>y = \dots</math> or <math>f^{-1}(x) = \dots</math></p> <p>or <math>-1 \leq (x - 1)/2 \leq 1</math>  must be '<math>x</math>', not <math>y</math> or <math>f(x)</math></p>	<p>at least one step attempted, or reasonable attempt at flow chart inversion</p> <p>(or any other variable provided same used on each side)</p> <p>condone '&lt;'s for M1  allow unsupported correct answers; <math>-1</math> to <math>3</math> is M1 A0</p>
<p><b>7</b> (A) True , (B) True , (C) False  Counterexample, e.g. <math>\sqrt{2} + (-\sqrt{2}) = 0</math></p>	<p>B2,1,0 B1 [3]</p>		

<b>8(i)</b> When $x = 1$ , $y = 3 \ln 1 + 1 - 1^2 = 0$	E1 [1]		
<b>(ii)</b> $\frac{dy}{dx} = \frac{3}{x} + 1 - 2x$ $\Rightarrow$ At R, $\frac{dy}{dx} = 0 = \frac{3}{x} + 1 - 2x$ $\Rightarrow 3 + x - 2x^2 = 0$ $\Rightarrow (3 - 2x)(1 + x) = 0$ $\Rightarrow x = 1.5$ , (or $-1$ ) $\Rightarrow y = 3 \ln 1.5 + 1.5 - 1.5^2 = 0.466$ (3 s.f.) $\frac{d^2y}{dx^2} = -\frac{3}{x^2} - 2$ When $x = 1.5$ , $d^2y/dx^2 (= -10/3) < 0 \Rightarrow \max$	M1 A1cao  M1 M1 A1 M1 A1cao  B1ft  E1 [9]	$d/dx (\ln x) = 1/x$  re-arranging into a quadratic = 0 factorising or formula or completing square substituting their $x$  ft their $dy/dx$ on equivalent work  www – don't need to calculate 10/3	SC1 for $x = 1.5$ unsupported, SC3 if verified    but condone rounding errors on 0.466
<b>(iii)</b> Let $u = \ln x$ , $du/dx = 1/x$ $dv/dx = 1$ , $v = x$ $\Rightarrow \int \ln x dx = x \ln x - \int x \cdot \frac{1}{x} dx$ $= x \ln x - \int 1 dx$ $= x \ln x - x + c$  $\Rightarrow A = \int_1^{2.05} (3 \ln x + x - x^2) dx$ $= \left[ 3x \ln x - 3x + \frac{1}{2}x^2 - \frac{1}{3}x^3 \right]_1^{2.05}$ $= -2.5057 + 2.833..$ $= 0.33$ (2 s.f.)	M1  A1  A1  B1  B1ft  M1dep A1cao [7]	parts   condone no $c$  correct integral and limits (soi)  $\left[ 3 \times \text{their } 'x \ln x - x' + \frac{1}{2}x^2 - \frac{1}{3}x^3 \right]$ substituting correct limits dep 1 <sup>st</sup> B1	allow correct result to be quoted (SC3)

<b>9(i)</b> $(0, \frac{1}{2})$	B1 [1]	allow $y = \frac{1}{2}$ , but not $(x =) \frac{1}{2}$ or $(\frac{1}{2}, 0)$ nor $P = 1/2$	
<b>(ii)</b> $\frac{dy}{dx} = \frac{(1+e^{2x})2e^{2x} - e^{2x} \cdot 2e^{2x}}{(1+e^{2x})^2}$ $= \frac{2e^{2x}}{(1+e^{2x})^2}$ When $x = 0$ , $dy/dx = 2e^0/(1+e^0)^2 = \frac{1}{2}$	M1 A1 A1  B1ft [4]	Quotient or product rule correct expression – condone missing bracket cao – mark final answer  follow through their derivative	product rule: $\frac{dy}{dx} = e^{2x} \cdot 2e^{2x}(-1)(1+e^{2x})^{-2} + 2e^{2x}(1+e^{2x})^{-1}$ $-\frac{2e^{2x}}{(1+e^{2x})^2}$ from $(udv - vdu)/v^2$ SC1
<b>(iii)</b> $A = \int_0^1 \frac{e^{2x}}{1+e^{2x}} dx$ $= \left[ \frac{1}{2} \ln(1+e^{2x}) \right]_0^1$ or let $u = 1 + e^{2x}$ , $du/dx = 2e^{2x}$ $\Rightarrow A = \int_2^{1+e^2} \frac{1/2}{u} du = \left[ \frac{1}{2} \ln u \right]_2^{1+e^2}$ $= \frac{1}{2} \ln(1+e^2) - \frac{1}{2} \ln 2$ $= \frac{1}{2} \ln \left[ \frac{1+e^2}{2} \right]^*$	B1  M1 A1  M1  A1  M1  E1 [5]	correct integral and limits (soi)  $k \ln(1+e^{2x})$ $k = \frac{1}{2}$  or $v = e^{2x}$ , $dv/dx = 2e^{2x}$ o.e.  [ $\frac{1}{2} \ln u$ ] or [ $\frac{1}{2} \ln(v+1)$ ]  substituting correct limits  www	condone no dx       allow missing dx's or incompatible limits, but penalise missing brackets
<b>(iv)</b> $g(-x) = \frac{1}{2} \left[ \frac{e^{-x} - e^x}{e^{-x} + e^x} \right] = -\frac{1}{2} \left[ \frac{e^x - e^{-x}}{e^x + e^{-x}} \right] = -g(x)$ Rotational symmetry of order 2 about O	M1 E1  B1 [3]	substituting $-x$ for $x$ in $g(x)$ completion www – taking out $-ve$ must be clear must have ‘rotational’ ‘about O’, ‘order 2’ (oe)	not $g(-x) \neq g(x)$ . Condone use of f for g.
<b>(v)(A)</b> $g(x) + \frac{1}{2} = \frac{1}{2} \cdot \frac{e^x - e^{-x}}{e^x + e^{-x}} + \frac{1}{2} = \frac{1}{2} \cdot \left( \frac{e^x - e^{-x} + e^x + e^{-x}}{e^x + e^{-x}} \right)$ $= \frac{1}{2} \cdot \left( \frac{2e^x}{e^x + e^{-x}} \right)$ $= \frac{e^x \cdot e^x}{e^x(e^x + e^{-x})} = \frac{e^{2x}}{e^{2x} + 1} = f(x)$ <b>(B)</b> Translation $\begin{pmatrix} 0 \\ 1/2 \end{pmatrix}$ <b>(C)</b> Rotational symmetry [of order 2] about P	M1 A1  E1 M1 A1  B1 [6]	combining fractions (correctly)    translation in y direction up $\frac{1}{2}$ unit dep ‘translation’ used o.e. condone omission of $180^\circ$ /order 2	allow ‘shift’, ‘move’ in correct direction for M1. $\begin{pmatrix} 0 \\ 1/2 \end{pmatrix}$ alone is SC1.

**OCR (Oxford Cambridge and RSA Examinations)**  
**1 Hills Road**  
**Cambridge**  
**CB1 2EU**

**OCR Customer Contact Centre**

**14 – 19 Qualifications (General)**

Telephone: 01223 553998

Facsimile: 01223 552627

Email: [general.qualifications@ocr.org.uk](mailto:general.qualifications@ocr.org.uk)

**[www.ocr.org.uk](http://www.ocr.org.uk)**

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

**Oxford Cambridge and RSA Examinations**  
**is a Company Limited by Guarantee**  
**Registered in England**  
**Registered Office; 1 Hills Road, Cambridge, CB1 2EU**  
**Registered Company Number: 3484466**  
**OCR is an exempt Charity**



**OCR (Oxford Cambridge and RSA Examinations)**  
**Head office**  
**Telephone: 01223 552552**  
**Facsimile: 01223 552553**

© OCR 2010