

**ADVANCED GCE  
MATHEMATICS**

**4727/01**

Further Pure Mathematics 3

**THURSDAY 24 JANUARY 2008**

Morning

Time: 1 hour 30 minutes

**Additional materials:** Answer Booklet (8 pages)  
List of Formulae (MF1)

**INSTRUCTIONS TO CANDIDATES**

- Write your name, centre number and candidate number in the spaces provided on the answer booklet.
- Read each question carefully and make sure you know what you have to do before starting your answer.
- Answer **all** the questions.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question or is clearly appropriate.
- You are permitted to use a graphical calculator in this paper.

**INFORMATION FOR CANDIDATES**

- The number of marks is given in brackets [ ] at the end of each question or part question.
- The total number of marks for this paper is 72.
- **You are reminded of the need for clear presentation in your answers.**

This document consists of 4 printed pages.

- 1 (a) A group  $G$  of order 6 has the combination table shown below.

	$e$	$a$	$b$	$p$	$q$	$r$
$e$	$e$	$a$	$b$	$p$	$q$	$r$
$a$	$a$	$b$	$e$	$r$	$p$	$q$
$b$	$b$	$e$	$a$	$q$	$r$	$p$
$p$	$p$	$q$	$r$	$e$	$a$	$b$
$q$	$q$	$r$	$p$	$b$	$e$	$a$
$r$	$r$	$p$	$q$	$a$	$b$	$e$

- (i) State, with a reason, whether or not  $G$  is commutative. [1]
- (ii) State the number of subgroups of  $G$  which are of order 2. [1]
- (iii) List the elements of the subgroup of  $G$  which is of order 3. [1]
- (b) A multiplicative group  $H$  of order 6 has elements  $e, c, c^2, c^3, c^4, c^5$ , where  $e$  is the identity. Write down the order of each of the elements  $c^3, c^4$  and  $c^5$ . [3]

- 2 Find the general solution of the differential equation

$$\frac{d^2y}{dx^2} - 8\frac{dy}{dx} + 16y = 4x. \quad [7]$$

- 3 Two fixed points,  $A$  and  $B$ , have position vectors  $\mathbf{a}$  and  $\mathbf{b}$  relative to the origin  $O$ , and a variable point  $P$  has position vector  $\mathbf{r}$ .
- (i) Give a geometrical description of the locus of  $P$  when  $\mathbf{r}$  satisfies the equation  $\mathbf{r} = \lambda\mathbf{a}$ , where  $0 \leq \lambda \leq 1$ . [2]
- (ii) Given that  $P$  is a point on the line  $AB$ , use a property of the vector product to explain why  $(\mathbf{r} - \mathbf{a}) \times (\mathbf{r} - \mathbf{b}) = \mathbf{0}$ . [2]
- (iii) Give a geometrical description of the locus of  $P$  when  $\mathbf{r}$  satisfies the equation  $\mathbf{r} \times (\mathbf{a} - \mathbf{b}) = \mathbf{0}$ . [3]

4 The integrals  $C$  and  $S$  are defined by

$$C = \int_0^{\frac{1}{2}\pi} e^{2x} \cos 3x \, dx \quad \text{and} \quad S = \int_0^{\frac{1}{2}\pi} e^{2x} \sin 3x \, dx.$$

By considering  $C + iS$  as a single integral, show that

$$C = -\frac{1}{13}(2 + 3e^\pi),$$

and obtain a similar expression for  $S$ .

[8]

(You may assume that the standard result for  $\int e^{kx} \, dx$  remains true when  $k$  is a complex constant, so that  $\int e^{(a+ib)x} \, dx = \frac{1}{a+ib} e^{(a+ib)x}$ .)

5 (i) Find the general solution of the differential equation

$$\frac{dy}{dx} + \frac{y}{x} = \sin 2x,$$

expressing  $y$  in terms of  $x$  in your answer.

[6]

In a particular case, it is given that  $y = \frac{2}{\pi}$  when  $x = \frac{1}{4}\pi$ .

(ii) Find the solution of the differential equation in this case.

[2]

(iii) Write down a function to which  $y$  approximates when  $x$  is large and positive.

[1]

6 A tetrahedron  $ABCD$  is such that  $AB$  is perpendicular to the base  $BCD$ . The coordinates of the points  $A$ ,  $C$  and  $D$  are  $(-1, -7, 2)$ ,  $(5, 0, 3)$  and  $(-1, 3, 3)$  respectively, and the equation of the plane  $BCD$  is  $x + 2y - 2z = -1$ .

(i) Find, in either order, the coordinates of  $B$  and the length of  $AB$ .

[5]

(ii) Find the acute angle between the planes  $ACD$  and  $BCD$ .

[6]

7 (i) (a) Verify, without using a calculator, that  $\theta = \frac{1}{8}\pi$  is a solution of the equation  $\sin 6\theta = \sin 2\theta$ .

[1]

(b) By sketching the graphs of  $y = \sin 6\theta$  and  $y = \sin 2\theta$  for  $0 \leq \theta \leq \frac{1}{2}\pi$ , or otherwise, find the other solution of the equation  $\sin 6\theta = \sin 2\theta$  in the interval  $0 < \theta < \frac{1}{2}\pi$ .

[2]

(ii) Use de Moivre's theorem to prove that

$$\sin 6\theta \equiv \sin 2\theta(16 \cos^4 \theta - 16 \cos^2 \theta + 3).$$

[5]

(iii) Hence show that one of the solutions obtained in part (i) satisfies  $\cos^2 \theta = \frac{1}{4}(2 - \sqrt{2})$ , and justify which solution it is.

[3]

8 Groups  $A$ ,  $B$ ,  $C$  and  $D$  are defined as follows:

$A$ : the set of numbers  $\{2, 4, 6, 8\}$  under multiplication modulo 10,

$B$ : the set of numbers  $\{1, 5, 7, 11\}$  under multiplication modulo 12,

$C$ : the set of numbers  $\{2^0, 2^1, 2^2, 2^3\}$  under multiplication modulo 15,

$D$ : the set of numbers  $\left\{\frac{1+2m}{1+2n}, \text{ where } m \text{ and } n \text{ are integers}\right\}$  under multiplication.

(i) Write down the identity element for each of groups  $A$ ,  $B$ ,  $C$  and  $D$ . [2]

(ii) Determine in each case whether the groups

$A$  and  $B$ ,

$B$  and  $C$ ,

$A$  and  $C$

are isomorphic or non-isomorphic. Give sufficient reasons for your answers. [5]

(iii) Prove the closure property for group  $D$ . [4]

(iv) Elements of the set  $\left\{\frac{1+2m}{1+2n}, \text{ where } m \text{ and } n \text{ are integers}\right\}$  are combined under **addition**. State which of the four basic group properties are **not** satisfied. (Justification is not required.) [2]