

**ADVANCED GCE
MATHEMATICS**

Probability & Statistics 3

TUESDAY 15 JANUARY 2008

4734/01

Morning
Time: 1 hour 30 minutes

Additional materials: Answer Booklet (8 pages)
List of Formulae (MF1)

INSTRUCTIONS TO CANDIDATES

- Write your name, centre number and candidate number in the spaces provided on the answer booklet.
- Read each question carefully and make sure you know what you have to do before starting your answer.
- Answer **all** the questions.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question or is clearly appropriate.
- You are permitted to use a graphical calculator in this paper.

INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [] at the end of each question or part question.
- The total number of marks for this paper is 72.
- **You are reminded of the need for clear presentation in your answers.**

This document consists of 4 printed pages.

- 1 A blueberry farmer increased the amount of water sprayed over his berries to see what effect this had on their weight. The farmer weighed each of a random sample of 80 berries of the previous season's crop and each of a random sample of 100 berries of the new crop. The results are summarised in the following table, in which \bar{x} denotes the sample mean weight in grams, and s^2 denotes an unbiased estimate of the relevant population variance.

	Sample size	\bar{x}	s^2
Previous season's crop (P)	80	1.24	0.003 56
New crop (N)	100	1.36	0.003 40

- (i) Calculate an estimate of $\text{Var}(\bar{X}_N - \bar{X}_P)$. [2]
- (ii) Calculate a 95% confidence interval for the difference in population mean weights. [3]
- (iii) Give a reason why it is unnecessary to use a t -distribution in calculating the confidence interval. [1]
- 2 The times taken for customers' phone complaints to be handled were monitored regularly by a company. During a particular week a researcher checked a random sample of 20 complaints and the times, x minutes, taken to handle the complaints are summarised by $\Sigma x = 337.5$. Handling times may be assumed to have a normal distribution with mean μ minutes and standard deviation 3.8 minutes.
- (i) Calculate a 98% confidence interval for μ . [4]
- During the same week two other researchers each calculated a 98% confidence interval for μ based on independent samples.
- (ii) Calculate the probability that at least one of the three intervals does not contain μ . [2]
- (iii) State two ways in which the calculation in part (i) would differ if the standard deviation were unknown. [2]
- 3 A transport authority wished to compare the performance of two rail companies, Western and Northern. They noted that the number of 'on-time' arrivals for a random sample of 80 Western trains over a particular route was 71. The corresponding number for a random sample of 90 Northern trains over a similar route was 73.
- (i) Test, at the 5% significance level, whether the population proportion of on-time Western trains exceeds the population proportion of on-time Northern trains. [7]
- (ii) Ranjit wishes to test whether the population proportion of on-time Western trains exceeds the population proportion of on-time Northern trains by more than 0.01. What variance estimate should she use? [2]

4 Eezimix flour is sold in small bags of weight S grams, where $S \sim N(502.1, 0.31^2)$. It is also sold in large bags of weight L grams, where $L \sim N(1004.9, 0.58^2)$.

(i) Find the probability that a randomly chosen large bag weighs at least 1 gram more than two randomly chosen small bags. [6]

(ii) Find the probability that a randomly chosen large bag weighs less than twice the weight of a randomly chosen small bag. [5]

5 Of two brands of lawnmower, A and B , brand A was claimed to take less time, on average, than brand B to mow similar stretches of lawn. In order to test this claim, 9 randomly selected gardeners were each given the task of mowing two regions of lawn, one with each brand of mower. All the regions had the same size and shape and had grass of the same height. The times taken, in seconds, are given in the table.

Gardener	1	2	3	4	5	6	7	8	9
Brand A	412	386	389	401	396	394	397	411	391
Brand B	422	394	385	408	394	399	397	410	397

(i) Test the claim using a paired-sample t -test at the 5% significance level. State a distributional assumption required for the test to be valid. [10]

(ii) Give a reason why a paired-sample t -test should be used, rather than a 2-sample t -test, in this case. [1]

[Questions 6 and 7 are printed overleaf.]

- 6 The Research and Development department of a paint manufacturer has produced paint of three different shades of grey, G_1 , G_2 and G_3 . In order to find the reaction of the public to these shades, each of a random sample of 120 people was asked to state which shade they preferred. The results, classified by gender, are shown in Table 1.

		Shade		
		G_1	G_2	G_3
Gender	Male	11	24	23
	Female	18	13	31

Table 1

Table 2 shows the corresponding expected values, correct to 2 decimal places, for a test of independence.

		Shade		
		G_1	G_2	G_3
Gender	Male	14.02	17.88	26.10
	Female	14.98	19.12	27.90

Table 2

- (i) Show how the value 17.88 for Male, G_2 was obtained. [2]
- (ii) Test, at the 5% significance level, whether gender and preferred shade are independent. [7]
- (iii) Determine the smallest significance level obtained from tables or calculator for which there is evidence that not all shades are equally preferred by people in general, irrespective of gender. [6]
- 7 The continuous random variable T has probability density function given by

$$f(t) = \begin{cases} 4t^3 & 0 < t \leq 1, \\ 0 & \text{otherwise.} \end{cases}$$

- (i) Obtain the cumulative distribution function of T . [2]
- (ii) Find the cumulative distribution function of H , where $H = \frac{1}{T^4}$, and hence show that the probability density function of H is given by $g(h) = \frac{1}{h^2}$ over an interval to be stated. [7]
- (iii) Find $E(1 + 2H^{-1})$. [3]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (OCR) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

OCR is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.