Centre Number			Candidate Number		
Surname					
Other Names					
Candidate Signature					



General Certificate of Education Advanced Level Examination January 2013

# **Mathematics**

MFP2

**Unit Further Pure 2** 

Wednesday 23 January 2013 9.00 am to 10.30 am

#### For this paper you must have:

• the blue AQA booklet of formulae and statistical tables. You may use a graphics calculator.

#### Time allowed

• 1 hour 30 minutes

## Instructions


- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer all questions.
- Write the question part reference (eg (a), (b)(i) etc) in the left-hand margin.
- You must answer each question in the space provided for that question. If you require extra space, use an AQA supplementary answer book; do **not** use the space provided for a different question.
- Do not write outside the box around each page.
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work that you do not want to be marked.

## Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 75.

#### **Advice**

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- You do not necessarily need to use all the space provided.



## Answer all questions.

Answer each question in the space provided for that question.		
1 (a)	Show that	
	$12\cosh x - 4\sinh x = 4e^x + 8e^{-x}$	(2 marks)
(b)	Solve the equation	
	$12\cosh x - 4\sinh x = 33$	
	giving your answers in the form $k \ln 2$ .	(5 marks)
QUESTION PART REFERENCE	Answer space for question 1	



QUESTION PART REFERENCE	Answer space for question 1



Two loci,  $L_1$  and  $L_2$ , in an Argand diagram are given by

$$L_1: |z + 6 - 5i| = 4\sqrt{2}$$

$$L_2: \quad \arg(z+i) = \frac{3\pi}{4}$$

The point P represents the complex number -2 + i.

- (a) Verify that the point P is a point of intersection of  $L_1$  and  $L_2$ . (2 marks)
- (b) Sketch  $L_1$  and  $L_2$  on one Argand diagram. (6 marks)
- (c) The point Q is also a point of intersection of  $L_1$  and  $L_2$ . Find the complex number that is represented by Q. (2 marks)

QUESTION PART REFERENCE	Answer space for question 2
·····	



QUESTION PART REFERENCE	Answer space for question 2
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
••••••	
••••••	
••••••	
•••••	
••••••	
••••••	
••••••	



- 3 (a) Show that  $\frac{1}{5r-2} \frac{1}{5r+3} = \frac{A}{(5r-2)(5r+3)}$ , stating the value of the constant A. (2 marks)
  - **(b)** Hence use the method of differences to show that

$$\sum_{r=1}^{n} \frac{1}{(5r-2)(5r+3)} = \frac{n}{3(5n+3)}$$
 (4 marks)

(c) Find the value of

$$\sum_{r=1}^{\infty} \frac{1}{(5r-2)(5r+3)}$$
 (1 mark)

Answer space for question 3	



QUESTION PART REFERENCE	Answer space for question 3
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
••••••	
••••••	
••••••	
••••••	
•••••	
••••••	
•••••	



4 The roots of the equation

$$z^3 - 5z^2 + kz - 4 = 0$$

are  $\alpha$ ,  $\beta$  and  $\gamma$ .

- (a) (i) Write down the value of  $\alpha + \beta + \gamma$  and the value of  $\alpha \beta \gamma$ . (2 marks)
  - (ii) Hence find the value of  $\alpha^2 \beta \gamma + \alpha \beta^2 \gamma + \alpha \beta \gamma^2$ . (2 marks)
- **(b)** The value of  $\alpha^2 \beta^2 + \beta^2 \gamma^2 + \gamma^2 \alpha^2$  is -4.
  - (i) Explain why  $\alpha$ ,  $\beta$  and  $\gamma$  cannot all be real. (1 mark)
  - (ii) By considering  $(\alpha\beta + \beta\gamma + \gamma\alpha)^2$ , find the possible values of k. (4 marks)

QUESTION PART REFERENCE	Answer space for question 4
•••••	
•••••	



QUESTION PART REFERENCE	Answer space for question 4



5 (a) Using the definition 
$$\tanh y = \frac{e^y - e^{-y}}{e^y + e^{-y}}$$
, show that, for  $|x| < 1$ ,

$$\tanh^{-1} x = \frac{1}{2} \ln \left( \frac{1+x}{1-x} \right) \tag{3 marks}$$

**(b)** Hence, or otherwise, show that 
$$\frac{d}{dx}(\tanh^{-1}x) = \frac{1}{1-x^2}$$
. (3 marks)

(c) Use integration by parts to show that

$$\int_0^{\frac{1}{2}} 4 \tanh^{-1} x \, \mathrm{d}x = \ln \left( \frac{3^m}{2^n} \right)$$

where m and n are positive integers.

(5 marks)

Answer space for question 5	
	••
	••



QUESTION PART REFERENCE	Answer space for question 5
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
••••••	
••••••	
••••••	
•••••	
•••••	
•••••	
•••••	
•••••	
••••••	
•••••	
• • • • • • • • • • • • • • • • • • • •	



QUESTION PART REFERENCE	Answer space for question 5
•••••	
•••••	
•••••	
••••••	



QUESTION PART REFERENCE	Answer space for question 5
•••••	
•••••	



**6** A curve is defined parametrically by

$$x = t^3 + 5$$
,  $y = 6t^2 - 1$ 

The arc length between the points where t = 0 and t = 3 on the curve is s.

- (a) Show that  $s = \int_0^3 3t \sqrt{t^2 + A} \, dt$ , stating the value of the constant A. (4 marks)
- (b) Hence show that s = 61. (4 marks)

QUESTION PART REFERENCE	Answer space for question 6



QUESTION PART REFERENCE	Answer space for question 6
••••••	
•••••	
•••••	
•••••	
•••••	
•••••	
••••••	
•••••	
•••••	
•••••	
•••••	
•••••	
••••••	
• • • • • • • • • • • • • • • • • • • •	



7		The polynomial $p(n)$ is given by $p(n) = (n-1)^3 + n^3 + (n+1)^3$ .	
(a	) (i)	Show that $p(k + 1) - p(k)$ , where k is a positive integer, is a multiple of 9.	(3 marks)
	(ii)	Prove by induction that $p(n)$ is a multiple of 9 for all integers $n \ge 1$ .	(4 marks)
(b	)	Using the result from part (a)(ii), show that $n(n^2 + 2)$ is a multiple of 3 for positive integer $n$ .	any (2 marks)
QUESTION PART REFERENCE	Ans	swer space for question 7	
	•••••		
	•••••		•••••
	••••••		•••••
	•••••		•••••
• • • • • • • • • • • • • • • • • • • •	•••••		•••••
	••••••		
	•••••		
	•••••		
	•••••		
	•••••		
	•••••		
	•••••		•••••
	••••••		
	•••••		



QUESTION PART REFERENCE	Answer space for question 7



- **8 (a)** Express  $-4 + 4\sqrt{3}i$  in the form  $re^{i\theta}$ , where r > 0 and  $-\pi < \theta \le \pi$ . (3 marks)
  - (b) (i) Solve the equation  $z^3 = -4 + 4\sqrt{3}i$ , giving your answers in the form  $re^{i\theta}$ , where r > 0 and  $-\pi < \theta \le \pi$ .
    - (ii) The roots of the equation  $z^3 = -4 + 4\sqrt{3}i$  are represented by the points P, Q and R on an Argand diagram.

Find the area of the triangle PQR, giving your answer in the form  $k\sqrt{3}$ , where k is an integer. (3 marks)

(c) By considering the roots of the equation  $z^3 = -4 + 4\sqrt{3}i$ , show that

$$\cos\frac{2\pi}{9} + \cos\frac{4\pi}{9} + \cos\frac{8\pi}{9} = 0 \tag{4 marks}$$

QUESTION PART REFERENCE	Answer space for question 8



QUESTION PART REFERENCE	Answer space for question 8
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
••••••	
•••••	
•••••	
••••••	
•••••	
•••••	
•••••	
••••••	
•••••	
••••••	
••••••	
•••••	
••••••	
•••••	



QUESTION PART REFERENCE	Answer space for question 8
•••••	
<u> </u>	
	END OF QUESTIONS
Copyria	ht © 2013 AQA and its licensors. All rights reserved.

