

General Certificate of Education
January 2006
Advanced Level Examination



MATHEMATICS
Unit Further Pure 4

MFP4

Monday 23 January 2006 1.30 pm to 3.00 pm

For this paper you must have:

- an 8-page answer book
- the **blue** AQA booklet of formulae and statistical tables

You may use a graphics calculator.

Time allowed: 1 hour 30 minutes

Instructions

- Use blue or black ink or ball-point pen. Pencil should only be used for drawing.
- Write the information required on the front of your answer book. The *Examining Body* for this paper is AQA. The *Paper Reference* is MFP4.
- Answer **all** questions.
- All necessary working should be shown; otherwise marks for method may be lost.

Information

- The maximum mark for this paper is 75.
- The marks for questions are shown in brackets.

Advice

- Unless stated otherwise, formulae may be quoted, without proof, from the booklet.

Answer **all** questions.

1 Describe the geometrical transformation defined by the matrix

$$\begin{bmatrix} 0.6 & 0 & 0.8 \\ 0 & 1 & 0 \\ -0.8 & 0 & 0.6 \end{bmatrix} \quad (3 \text{ marks})$$

2 The matrices \mathbf{P} and \mathbf{Q} are defined in terms of the constant k by

$$\mathbf{P} = \begin{bmatrix} 3 & 2 & 1 \\ 1 & -1 & k \\ 5 & 3 & 2 \end{bmatrix} \quad \text{and} \quad \mathbf{Q} = \begin{bmatrix} 5 & 4 & 1 \\ 3 & k & -1 \\ 7 & 3 & 2 \end{bmatrix}$$

(a) Express $\det \mathbf{P}$ and $\det \mathbf{Q}$ in terms of k . (3 marks)

(b) Given that $\det(\mathbf{PQ}) = 16$, find the two possible values of k . (4 marks)

3 (a) The plane Π has equation $\mathbf{r} = \begin{bmatrix} 2 \\ 5 \\ 1 \end{bmatrix} + \lambda \begin{bmatrix} 0 \\ 3 \\ 1 \end{bmatrix} + \mu \begin{bmatrix} 4 \\ -1 \\ 0 \end{bmatrix}$.

(i) Find a vector which is perpendicular to both $\begin{bmatrix} 0 \\ 3 \\ 1 \end{bmatrix}$ and $\begin{bmatrix} 4 \\ -1 \\ 0 \end{bmatrix}$. (2 marks)

(ii) Hence find an equation for Π in the form $\mathbf{r} \cdot \mathbf{n} = d$. (2 marks)

(b) The line L has equation $\left(\mathbf{r} - \begin{bmatrix} -1 \\ 2 \\ 6 \end{bmatrix} \right) \times \begin{bmatrix} 3 \\ 0 \\ -1 \end{bmatrix} = \mathbf{0}$.

Verify that $\mathbf{r} = \begin{bmatrix} 2 \\ 2 \\ 5 \end{bmatrix} + t \begin{bmatrix} 3 \\ 0 \\ -1 \end{bmatrix}$ is also an equation for L . (2 marks)

(c) Determine the position vector of the point of intersection of Π and L . (4 marks)

4 The vectors \mathbf{a} , \mathbf{b} and \mathbf{c} are given by

$$\mathbf{a} = \mathbf{i} - \mathbf{j} - \mathbf{k}, \quad \mathbf{b} = 2\mathbf{i} + 3\mathbf{j} - \mathbf{k} \quad \text{and} \quad \mathbf{c} = 4\mathbf{i} - \mathbf{j} + 5\mathbf{k}$$

(a) (i) Evaluate $\begin{vmatrix} 1 & -1 & -1 \\ 2 & 3 & -1 \\ 4 & -1 & 5 \end{vmatrix}$. (2 marks)

(ii) Hence determine whether \mathbf{a} , \mathbf{b} and \mathbf{c} are linearly dependent or independent. (1 mark)

(b) (i) Evaluate $\mathbf{b} \cdot \mathbf{c}$. (2 marks)

(ii) Show that $\mathbf{b} \times \mathbf{c}$ can be expressed in the form $m\mathbf{a}$, where m is a scalar. (2 marks)

(iii) Use these results to describe the geometrical relationship between \mathbf{a} , \mathbf{b} and \mathbf{c} . (1 mark)

(c) The points A , B and C have position vectors \mathbf{a} , \mathbf{b} and \mathbf{c} respectively relative to an origin O . The points O , A , B and C are four of the eight vertices of a cuboid. Determine the volume of this cuboid. (2 marks)

5 The transformation T maps (x, y) to (x', y') , where

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

(a) Describe the difference between *an invariant line* and *a line of invariant points* of T . (1 mark)

(b) Evaluate the determinant of the matrix $\begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix}$ and describe the geometrical significance of the result in relation to T . (2 marks)

(c) Show that T has a line of invariant points, and find a cartesian equation for this line. (2 marks)

(d) (i) Find the image of the point $(x, -x + c)$ under T . (2 marks)

(ii) Hence show that all lines of the form $y = -x + c$, where c is an arbitrary constant, are invariant lines of T . (2 marks)

(e) Describe the transformation T geometrically. (3 marks)

Turn over ►

6 (a) Show that

$$\begin{vmatrix} 1 & 1 & 1 \\ a & b & c \\ bc & ca & ab \end{vmatrix} = (a-b)(b-c)(c-a) \quad (5 \text{ marks})$$

(b) (i) Hence, or otherwise, show that the system of equations

$$\begin{aligned} x + y + z &= p \\ 3x + 3y + 5z &= q \\ 15x + 15y + 9z &= r \end{aligned}$$

has no unique solution whatever the values of p , q and r . (2 marks)

(ii) Verify that this system is consistent when $24p - 3q - r = 0$. (2 marks)

(iii) Find the solution of the system in the case where $p = 1$, $q = 8$ and $r = 0$. (5 marks)

7 The matrix $\mathbf{M} = \begin{bmatrix} 1 & -1 & 1 \\ 3 & -3 & 1 \\ 3 & -5 & 3 \end{bmatrix}$.

(a) Given that $\mathbf{u} = \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}$ and $\mathbf{v} = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$ are eigenvectors of \mathbf{M} , find the eigenvalues corresponding to \mathbf{u} and \mathbf{v} . (5 marks)

(b) Given also that the third eigenvalue of \mathbf{M} is 1, find a corresponding eigenvector. (6 marks)

(c) (i) Express the vector $\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$ in terms of \mathbf{u} and \mathbf{v} . (1 mark)

(ii) Deduce that $\mathbf{M}^n \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = \lambda^n \mathbf{u} + \mu^n \mathbf{v}$, where λ and μ are scalar constants whose values should be stated. (4 marks)

(iii) Hence prove that, for all positive **odd** integers n ,

$$\mathbf{M}^n \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = \begin{bmatrix} 2^n \\ 0 \\ 2^n \end{bmatrix} \quad (3 \text{ marks})$$

END OF QUESTIONS

There are no questions printed on this page

There are no questions printed on this page

There are no questions printed on this page