Centre Number			Candidate Number		
Surname					
Other Names					
Candidate Signature					



General Certificate of Education Advanced Level Examination June 2010

Mathematics

MFP4

Unit Further Pure 4

Tuesday 15 June 2010 9.00 am to 10.30 am

For this paper you must have:

• the blue AQA booklet of formulae and statistical tables. You may use a graphics calculator.

Time allowed

• 1 hour 30 minutes

Instructions


- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer all questions.
- Write the question part reference (eg (a), (b)(i) etc) in the left-hand margin.
- You must answer the questions in the spaces provided. Do not write outside the box around each page.
- Show all necessary working; otherwise marks for method may be lost
- Do all rough work in this book. Cross through any work that you do not want to be marked.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 75.

Advice

 Unless stated otherwise, you may quote formulae, without proof, from the booklet.



Answer all questions in the spaces provided.

1 The position vectors of the points P, Q and R are, respectively,

$$\mathbf{p} = \begin{bmatrix} 3 \\ 4 \\ -1 \end{bmatrix}, \quad \mathbf{q} = \begin{bmatrix} -1 \\ 2 \\ 2 \end{bmatrix} \quad \text{and} \quad \mathbf{r} = \begin{bmatrix} 1 \\ 4 \\ 1 \end{bmatrix}$$

(a) Show that p, q and r are linearly dependent.

(2 marks)

(b) Determine the area of triangle PQR.

(4 marks)

QUESTION PART REFERENCE	
•••••	
•••••	



QUESTION PART REFERENCE	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	



(2 marks)

2 Let $A =$	$\begin{bmatrix} 1 \\ 2 \end{bmatrix}$	$\begin{bmatrix} x \\ 3 \end{bmatrix}$, B =	$\begin{bmatrix} 1 \\ 2 \end{bmatrix}$	$\begin{bmatrix} -1 \\ 2 \end{bmatrix}$	and $C =$	$\begin{bmatrix} 4 - 4x \\ 8x - 4 \end{bmatrix}$	8 4	
-------------	--	--	--------------	--	---	-----------	--	-----	--

- (a) Find AB in terms of x.
- (b) Show that $\mathbf{B}^{\mathrm{T}}\mathbf{A}^{\mathrm{T}} = \mathbf{C}$ for some value of x. (5 marks)

QUESTION	
PART	
REFERENCE	
• • • • • • • • • •	
• • • • • • • • • •	



QUESTION PART REFERENCE	
•••••	
••••••	
••••••	
••••••	
••••••	
••••••	
•••••	
••••••	
•••••	



3	The plane Π_1 is perpendicular to the vector $9\mathbf{i} - 8\mathbf{j} + 72\mathbf{k}$ and passes through the point $A(2, 10, 1)$.
(a	Find, in the form $\mathbf{r} \cdot \mathbf{n} = d$, a vector equation for Π_1 . (3 marks)
(b	Determine the exact value of the cosine of the acute angle between Π_1 and the plane Π_2 with equation $\mathbf{r} \cdot (\mathbf{i} + \mathbf{j} + \mathbf{k}) = 11$. (4 marks)
QUESTION PART REFERENCE	
••••••	
•••••	
••••••	
•••••	
•••••	
•••••	



QUESTION PART REFERENCE	
•••••	
•••••	
•••••	
•••••	
•••••	



4 The fixed points A and B and the variable point C have position vectors

$$\mathbf{a} = \begin{bmatrix} 3 \\ -4 \\ 1 \end{bmatrix}, \quad \mathbf{b} = \begin{bmatrix} 2 \\ 1 \\ -3 \end{bmatrix} \quad \text{and} \quad \mathbf{c} = \begin{bmatrix} 2 - t \\ t \\ 5 \end{bmatrix}$$

respectively, relative to the origin O, where t is a scalar parameter.

- (a) Find an equation of the line AB in the form $(\mathbf{r} \mathbf{u}) \times \mathbf{v} = \mathbf{0}$. (3 marks)
- (b) Determine $\mathbf{b} \times \mathbf{c}$ in terms of t. (4 marks)
- (c) (i) Show that $\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})$ is constant for all values of t, and state the value of this constant. (2 marks)
 - (ii) Write down a geometrical conclusion that can be deduced from the answer to part (c)(i). (1 mark)

QUESTION PART REFERENCE	
PART	
REFERENCE	
	•
• • • • • • • • • • • • • • • • • • • •	



QUESTION PART REFERENCE	
•••••	
••••••	
••••••	
••••••	
••••••	
••••••	
•••••	
••••••	
•••••	



5	Factorise fully the determinant	$\begin{vmatrix} x \\ x^2 \\ yz \end{vmatrix}$	y y^2 zx	$\begin{bmatrix} z \\ z^2 \\ xy \end{bmatrix}$		(8 marks)
QUESTION PART						
REFERENCE						
		•••••	•••••	• • • • • • •		
•••••		•••••	•••••	•••••	••••••	
		•••••	•••••	•••••		
•••••		•••••	•••••	•••••	••••••	
		•••••	•••••	• • • • • • •		
		•••••	•••••	•••••	••••••	
				•••••		
		•••••	•••••	•••••		
•••••		•••••	•••••	•••••	••••••	
					•••••	
		•••••	•••••	•••••		
		•••••	•••••	•••••	•••••	
		•••••	•••••	• • • • • •		
•••••		•••••	•••••	•••••	••••••	
		•••••	•••••	• • • • • • •		
•••••		•••••	•••••	•••••	•••••••••••	
		•••••	•••••	•••••		
• • • • • • • • • • • • • • • • • • • •		•••••	•••••	• • • • • • •	•••••	



QUESTION PART REFERENCE	
•••••	
•••••	
••••••	
•••••	
•••••	
•••••	
•••••	



6 The line L and the plane Π have vector equations

$$\mathbf{r} = \begin{bmatrix} 7 \\ 8 \\ 50 \end{bmatrix} + t \begin{bmatrix} 6 \\ 2 \\ -9 \end{bmatrix} \quad \text{and} \quad \mathbf{r} = \begin{bmatrix} -2 \\ 0 \\ -25 \end{bmatrix} + \lambda \begin{bmatrix} 5 \\ 3 \\ 4 \end{bmatrix} + \mu \begin{bmatrix} 1 \\ 6 \\ 2 \end{bmatrix}$$

respectively.

(a) (i) Find direction cosines for L.

(2 marks)

(ii) Show that L is perpendicular to Π .

(3 marks)

(b) For the system of equations

$$6p + 5q + r = 9$$

$$2p + 3q + 6r = 8$$

$$-9p + 4q + 2r = 75$$

form a pair of equations in p and q only, and hence find the unique solution of this system of equations. (5 marks)

- (c) It is given that L meets Π at the point P.
 - (i) Demonstrate how the coordinates of P may be obtained from the system of equations in part (b). (2 marks)
 - (ii) Hence determine the coordinates of P.

(2 marks)

PART REFERENCE	
••••••	
••••••	
•••••	

QUESTION PART REFERENCE	
•••••	



QUESTION PART REFERENCE	
••••••	
•••••	
•••••	
••••••	
••••••	
•••••	
•••••	
•••••	
••••••	



QUESTION PART REFERENCE	
••••••	
•••••	
•••••	
••••••	
••••••	
•••••	
•••••	
•••••	
••••••	



7 The transformation T is represented by the matrix **M** with diagonalised form

$$\mathbf{M} = \mathbf{U} \, \mathbf{D} \, \mathbf{U}^{-1}$$

where
$$\mathbf{U} = \begin{bmatrix} 4 & -1 \\ 1 & 3 \end{bmatrix}$$
 and $\mathbf{D} = \begin{bmatrix} 27 & 0 \\ 0 & 1 \end{bmatrix}$.

- (a) (i) State the eigenvalues, and corresponding eigenvectors, of M. (4 marks)
 - (ii) Find a cartesian equation for the line of invariant points of T. (2 marks)
- (b) Write down U^{-1} , and hence find the matrix M in the form

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

where a, b, c and d are integers.

(5 marks)

(c) By finding the element in the first row, first column position of \mathbf{M}^n , prove that

$$4 \times 3^{3n+1} + 1$$

is a multiple of 13 for all positive integers n.

(5 marks)

OUESTION PART REFERENCE	QUESTION	
	PART	
	REFERENCE	
	• • • • • • • • • • • • • • • • • • • •	
	• • • • • • • • • •	
	•••••	
	• • • • • • • • • •	
	•••••	



QUESTION PART REFERENCE	
••••••	
•••••	
•••••	
••••••	
••••••	
•••••	
•••••	
•••••	
••••••	



QUESTION PART REFERENCE	
•••••	
•••••	
••••••	
••••••	
••••••	
•••••	
•••••	
••••••	
••••••	
••••••	
•••••	



QUESTION PART REFERENCE	
•••••	



8	The matrix $\begin{bmatrix} 12 & 16 \\ -9 & 36 \end{bmatrix}$ represents the transformation which is the composition, in either order, of the two plane transformations
	E: an enlargement, centre O and scale factor k $(k > 0)$
	and
	S: a shear parallel to the line l which passes through O
	Show that $k = 24$ and find a cartesian equation for l . (7 marks)
QUESTION PART REFERENCE	
••••••	
••••••	
•••••	
••••••	
•••••	
•••••	



QUESTION PART REFERENCE	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	



QUESTION PART REFERENCE	
REFERENCE	
•••••	
•••••	



QUESTION PART REFERENCE	
	L
	END OF QUESTIONS



