



General Certificate of Education

Mathematics 6360

MM1A Mechanics 1A

Mark Scheme

2008 examination - January series

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation meeting attended by all examiners and is the scheme which was used by them in this examination. The standardisation meeting ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for the standardisation meeting each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed at the meeting and legislated for. If, after this meeting, examiners encounter unusual answers which have not been discussed at the meeting they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available to download from the AQA Website: www.aqa.org.uk

Copyright © 2008 AQA and its licensors. All rights reserved.

COPYRIGHT

AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

Key to mark scheme and abbreviations used in marking

M	mark is for method		
m or dM	mark is dependent on one or more M marks and is for method		
A	mark is dependent on M or m marks and is for accuracy		
B	mark is independent of M or m marks and is for method and accuracy		
E	mark is for explanation		
✓ or ft or F	follow through from previous incorrect result	MC	mis-copy
CAO	correct answer only	MR	mis-read
CSO	correct solution only	RA	required accuracy
AWFW	anything which falls within	FW	further work
AWRT	anything which rounds to	ISW	ignore subsequent work
ACF	any correct form	FIW	from incorrect work
AG	answer given	BOD	given benefit of doubt
SC	special case	WR	work replaced by candidate
OE	or equivalent	FB	formulae book
A2,1	2 or 1 (or 0) accuracy marks	NOS	not on scheme
–x EE	deduct x marks for each error	G	graph
NMS	no method shown	c	candidate
PI	possibly implied	sf	significant figure(s)
SCA	substantially correct approach	dp	decimal place(s)

No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded. However, there are situations in some units where part marks would be appropriate, particularly when similar techniques are involved. Your Principal Examiner will alert you to these and details will be provided on the mark scheme.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.


Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

Otherwise we require evidence of a correct method for any marks to be awarded.

Q	Solution	Marks	Total	Comments
1. (a)	$8 = \frac{1}{2}a \times 5^2$ $a = \frac{2 \times 8}{25} = 0.64 \text{ ms}^{-2}$ AG	M1		Use of constant acceleration equation with $u = 0$ to find a .
		A1	2	Correct answer from correct working, showing evidence of solving for a . Allow verification / substitution.
(b)	$T - 70 \times 9.8 = 70 \times 0.64$ $T = 730.8 = 731 \text{ N to 3 sf}$	M1		Three term equation of motion for crate.
		A1 A1	3	Correct equation Correct tension
(c)	$v = \frac{8}{5} = 1.6 \text{ ms}^{-1}$	B1	1	Correct average speed. Accept $\frac{8}{5}$ Allow $\frac{3.2 + 0}{2} = 1.6 \text{ ms}^{-1}$
	Total		6	
2.(a)	$U = \sqrt{10^2 - 8^2} = 6$	M1		Expression/equation for U based on a right angled triangle.
		A1	2	Correct U . Note $10^2 + 8^2$ gives M1A0
(b)	$\cos \theta = \frac{8}{10}$ $\theta = 037^\circ$	M1		Use of trigonometry to find angle.
		A1	2	Allow $\left\{ \begin{array}{l} \tan \theta = \frac{8}{6} \text{ or } \frac{6}{8} \\ \sin / \cos \theta = \frac{8}{10} \text{ or } \frac{6}{10} \end{array} \right.$ Correct angle. Accept 36.9° etc. Note 143° gives M1A0
	Total		4	

MM1A (cont)

Q	Solution	Marks	Total	Comments
3.(a)		B1	1	Diagram with three forces, labels and arrow heads. Different variables must be used for each tension
(b)	$T_1 \sin 30^\circ = 4 \times 9.8$ $T_1 = \frac{4 \times 9.8}{\sin 30^\circ} = 78.4 \text{ N}$	M1 A1 A1	3	Two term equation from resolving vertically. Must see a sin or cos term for M1 Correct equation Correct tension from correct working.
(c)	$T_2 = 78.4 \cos 30^\circ = 67.9 \text{ N}$	M1 A1	2	Two term equation from resolving horizontally. Correct tension.
	Total		6	
4. (a)(i)	$5 \begin{bmatrix} 2U \\ U \end{bmatrix} + 15 \begin{bmatrix} V \\ -1 \end{bmatrix} = 20 \begin{bmatrix} V \\ 0 \end{bmatrix}$ $5U - 15 = 0$ $U = 3$	M1 dM1 A1	3	Three term equation for conservation of momentum. Equation for U based on conservation of momentum. Correct value for U . Deduct one mark for using weight instead of mass.
(a)(ii)	$30 + 15V = 20V$ $30 = 5V$ $V = \frac{30}{5} = 6$	M1 A1F	2	Equation for V based on conservation of momentum. Correct value for V . Deduct one mark for using weight instead of mass.
	Total		5	

MM1A (cont)

Q	Solution	Marks	Total	Comments
5.(a)	$R = 15 \times 9.8 = 147 \text{ N}$ $F = 0.2 \times 147 = 29.4 \text{ N}$	M1 M1 A1	3	Finding normal reaction. Use of $F = \mu R$ Correct friction force.
(b)	$5g - T = 5a$ $T - 29.4 = 15a$ AG $49 - 29.4 = 20a$ $a = 0.98 \text{ ms}^{-2}$	M1 A1 M1 A1 A1	5	Three term equation of motion for 5kg particle Correct equation Three term equation of motion for 15kg block Correct equation Correct final answer from correct working.
(c)	$1.4^2 = 0^2 + 2 \times 0.98s$ $s = \frac{1.4^2}{1.96} = 1 \text{ m}$	M1 A1 A1	3	Use of a constant acceleration equation with $u = 0$ Correct equation Correct distance
	Total		11	

MM1A (cont)

6.(a)	$-3 = 70 \sin 10^\circ t - 4.9t^2$	M1		Vertical equation to find t containing a 3
		A1		Correct RHS of equation
	$4.9t^2 - 70 \sin 10^\circ t - 3 = 0$	A1		Correct LHS of equation
	$t = \frac{70 \sin 10^\circ \pm \sqrt{(70 \sin 10^\circ)^2 - 4 \times 4.9 \times (-3)}}{2 \times 4.9}$	dM1		Solving for t
	$t = -0.226$ or 2.71	dM1		Selecting the positive t
	$t = 2.71$	A1	6	Correct time from correct working
	(b) $s = 70 \cos 10^\circ \times 2.71 = 187$	M1		Finding horizontal distance
		A1	2	Correct distance
	(c) $v_x = 70 \cos 10^\circ = 68.9$	M1		Calculating horizontal component of velocity
		A1	2	Correct value
	(d) $v_y = 70 \sin 10^\circ - 9.8 \times 2.71 = -14.4$	M1		Calculating vertical component of velocity at $t = 2.71$
		A1		Correct value
	$v = \sqrt{68.9^2 + 14.4^2} = 70.4$	dM1		Finding speed
		A1	4	Correct speed
	Total		14	

MM1A (cont)

Q	Solution	Marks	Total	Comments
7.(a)	$4\mathbf{i} = 5\mathbf{j} + 40\mathbf{a}$ $\mathbf{a} = \frac{4\mathbf{i} - 5\mathbf{j}}{40} = 0.1\mathbf{i} - 0.125\mathbf{j}$ AG	M1 A1 dM1 A1	4	Forming a vector equation based on constant acceleration Correct equation Solving for \mathbf{a} Correct \mathbf{a} from correct working For $\frac{4\mathbf{i} - 5\mathbf{j}}{40}$ on its own give M0 Allow verification
(b)	$\mathbf{r} = 5\mathbf{j} \times 40 + \frac{1}{2}(0.1\mathbf{i} - 0.125\mathbf{j}) \times 40^2$ $= 80\mathbf{i} + 100\mathbf{j}$	M1 A1 A1	3	Finding position vector Correct expression Correct simplified result
(c)(i)	$\mathbf{v} = 5\mathbf{j} + (0.1\mathbf{i} - 0.125\mathbf{j})t$ $= 0.1t\mathbf{i} + (5 - 0.125t)\mathbf{j}$ $5 - 0.125t = -0.1t$ $5 = 0.025t$ $t = \frac{5}{0.025} = 200$	M1 A1 dM1 A1 A1	5	Expression for \mathbf{v} Correct expression for \mathbf{v} seen or implied Equating components, with or without a minus sign Correct equation Correct time.
(c)(ii)	$\mathbf{v} = 0.1 \times 200\mathbf{i} + (5 - 0.125 \times 200)\mathbf{j}$ $= 20\mathbf{i} - 20\mathbf{j}$	M1 A1F	2	Finding velocity using their time Correct velocity for their time
Total			14	

Note for question 7. Consistent use of $\mathbf{u} = 4\mathbf{i}$ or $5\mathbf{i}$ or $\mathbf{a} = 0.1\mathbf{i} + 0.125\mathbf{j}$ award method marks only.