



General Certificate of Education (A-level)
June 2011

Mathematics

MPC1

(Specification 6360)

Pure Core 1

Final

Mark Scheme

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all examiners participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for standardisation each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, examiners encounter unusual answers which have not been raised they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available from: aqa.org.uk

Copyright © 2011 AQA and its licensors. All rights reserved.

Copyright

AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

Key to mark scheme abbreviations

| | |
|-------------|--|
| M | mark is for method |
| m or dM | mark is dependent on one or more M marks and is for method |
| A | mark is dependent on M or m marks and is for accuracy |
| B | mark is independent of M or m marks and is for method and accuracy |
| E | mark is for explanation |
| ✓or ft or F | follow through from previous incorrect result |
| CAO | correct answer only |
| CSO | correct solution only |
| AWFW | anything which falls within |
| AWRT | anything which rounds to |
| ACF | any correct form |
| AG | answer given |
| SC | special case |
| OE | or equivalent |
| A2,1 | 2 or 1 (or 0) accuracy marks |
| –x EE | deduct x marks for each error |
| NMS | no method shown |
| PI | possibly implied |
| SCA | substantially correct approach |
| c | candidate |
| sf | significant figure(s) |
| dp | decimal place(s) |

No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

Otherwise we require evidence of a correct method for any marks to be awarded.

MPC1

| Q | Solution | Marks | Total | Comments |
|-------------|---|-------|----------|---|
| 1(a) | $y = \frac{13}{3} - \frac{7}{3}x$ | M1 | 2 | attempt at $y = a + bx$ or $\frac{\Delta y}{\Delta x}$ with 2 correct points |
| | (gradient =) $-\frac{7}{3}$ | A1 | | condone slip in rearranging if gradient is correct |
| | (b)(i) $y - 3 = \text{'their grad'}(x - -1)$ | M1 | 2 | or $7x + 3y = k$ and attempt at k using $x = -1$ and $y = 3$ or $y = (\text{their } m)x + c$ and attempt at c using $x = -1$ and $y = 3$ |
| | $y - 3 = -\frac{7}{3}(x + 1)$ or $7x + 3y = 2$ or $y = -\frac{7}{3}x + c, \quad c = \frac{2}{3}$ | A1cso | | correct equation in any form and replacing $--$ with $+$ sign |
| | (ii) $(4, -5)$ | B1,B1 | 2 | $x = 4, y = -5$ withhold if clearly from incorrect working |
| (c) | $7x + 3y = 13$ and $3x + 2y = 12$ \Rightarrow equation in x or y only | M1 | 3 | must use correct pair of equations and attempt to eliminate y (or x) |
| | $x = -2$ | A1 | | |
| | $y = 9$ | A1 | | |
| | Total | | 9 | |


MPC1 (cont)

| Q | Solution | Marks | Total | Comments |
|--------------|---|-------|----------|--|
| 2(a)(i) | $\sqrt{48} = 4\sqrt{3}$ | B1 | 1 | condone $k = 4$ stated |
| (ii) | $\frac{4\sqrt{3} + 6\sqrt{3}}{2\sqrt{3}}$ | M1 | | attempt to write each term in form $k\sqrt{3}$ with at least 2 terms correctly obtained |
| | | A1 | | correct unsimplified in terms of $\sqrt{3}$ only |
| | $= 5$ | A1cso | 3 | must simplify fraction to 5 |
| | | | | Alternative 1 $\times \frac{\sqrt{12}}{\sqrt{12}} \left(\text{or } \times \frac{\sqrt{3}}{\sqrt{3}} \right)$ M1 correct with integer terms $= \frac{24 + 36}{12}$ A1 $= 5$ A1cso |
| | | | | Alternative 2 $\frac{\sqrt{48} + \sqrt{108}}{\sqrt{12}}$ M1 $= \sqrt{4} + \sqrt{9}$ A1 $= 5$ A1cso |
| | | | | Alternative 3 $\sqrt{\frac{48}{12}} + 2\sqrt{\frac{27}{12}}$ M1 $= 2 + 2\sqrt{\frac{9}{4}}$ A1 $= 5$ A1cso |
| | | | | if hybrid of methods used, award M1 and most appropriate first A1 |
| | | | | NMS (answer =) 5 scores full marks |
| (b) | $\frac{1 - 5\sqrt{5}}{3 + \sqrt{5}} \times \frac{3 - \sqrt{5}}{3 - \sqrt{5}}$ | M1 | | |
| | (numerator =) $3 - \sqrt{5} - 15\sqrt{5} + 25$ | m1 | | correct unsimplified but must write $5\sqrt{5}\sqrt{5} = 25$ PI by 28 seen later |
| | (denominator = $9 - 5$) 4 | B1 | | must be seen as denominator |
| | giving $\frac{28 - 16\sqrt{5}}{4}$ | | | |
| | (answer =) $7 - 4\sqrt{5}$ | A1 | 4 | $m = 7, n = -4$ |
| Total | | | 8 | |

MPC1 (cont)

| Q | Solution | Marks | Total | Comments |
|--------------|--|------------------------|-----------|--|
| 3(a) | $\left(\frac{dV}{dt} = \right) \frac{3t^2}{4} - 3$ | M1 A1 | 2 | one of these terms correct all correct (no + c etc) |
| (b)(i) | $t = 1 \Rightarrow \frac{dV}{dt} = \frac{3}{4} - 3$ $= -2\frac{1}{4}$ | M1 A1cso | 2 | substituting $t = 1$ into their $\frac{dV}{dt}$ (-2.25 OE) BUT must have $\frac{dV}{dt}$ correct |
| (ii) | Volume is decreasing when $t = 1$ because $\frac{dV}{dt} < 0$ | E1✓ | 1 | must have used $\frac{dV}{dt}$ in (b)(i) or starts again must state that $\frac{dV}{dt} < 0$ (or $-2\frac{1}{4} < 0$ etc) ft increasing plus explanation if their $\frac{dV}{dt} > 0$ |
| (c)(i) | $\left(\frac{dV}{dt} = 0 \Rightarrow \right) \frac{3t^2}{4} - 3 = 0$ $\Rightarrow t^2 = 4$ $t = 2$ | M1 A1✓ A1cso | 3 | PI by “correct” equation being solved obtaining $t'' = k$ correctly from their $\frac{dV}{dt}$ withhold if answer left as $t = \pm 2$ |
| (ii) | $\left(\frac{d^2V}{dt^2} = \right) \frac{3t}{2}$ When $t = 2$, $\frac{d^2V}{dt^2} = 3$ or $\frac{d^2V}{dt^2} > 0$ \Rightarrow minimum | B1✓ M1 A1cso | 3 | (condone unsimplified) ft their $\frac{dV}{dt}$ ft their $\frac{d^2V}{dt^2}$ and value of t from (c)(i) |
| Total | | | 11 | |


MPC1 (cont)

| Q | Solution | Marks | Total | Comments |
|---------------|--|-------|-----------|---|
| 4(a) | $(x+2.5)^2$ | B1 | 3 | $p = \frac{5}{2}$ |
| | $q = 7 - \text{'their' } p^2$ | M1 | | unsimplified attempt at $q = 7 - \text{'their' } p^2$ |
| | $(x+2.5)^2 + 0.75$ | A1 | | $q = 7 - \frac{25}{4} = \frac{3}{4}$ |
| | <i>mark their final line as their answer</i> | | | |
| | | | | |
| (b)(i) | $x = -\text{'their' } p \quad \text{or} \quad y = \text{'their' } q$ | M1 | 2 | or $x = -\frac{5}{2}$ cao found using calculus |
| | $\left(-\frac{5}{2}, \frac{3}{4}\right)$ | A1cao | | condone correct coordinates stated $x = -2.5, \quad y = 0.75$ |
| (ii) | $x = -\frac{5}{2}$ | B1✓ | 1 | correct or ft “ $x = -\text{'their' } p$ ” |
| (iii) |  | B1 | 3 | y intercept = 7 stated or seen in table as $y = 7$ when $x = 0$ or 7 marked as intercept on y-axis (any graph) |
| | | M1 | | ∪ shape |
| | | A1 | | vertex above x-axis in correct quadrant and parabola extending beyond y-axis into first quadrant |
| (c) | Translation through $\begin{bmatrix} -\frac{5}{2} \\ \frac{3}{4} \end{bmatrix}$ | E1 | 3 | and no other transformation |
| | | M1 | | ft either ‘their’ $-p$ or ‘their’ q or one component correct for M1 |
| | | A1cao | | both components correct for A1; may describe in words or use a vector |
| | Total | | 12 | |


MPC1 (cont)

| Q | Solution | Marks | Total | Comments |
|---------------|---|--|----------|---|
| 5(a) | $p(3) = 3^3 - 2 \times 3^2 + 3 (= 27 - 18 + 3)$ $= 12$ | M1 A1 | 2 | p(3) attempted; not long division |
| (b) | $p(-1) = (-1)^3 - 2(-1)^2 + 3$ $p(-1) = -1 - 2 + 3 = 0 \Rightarrow x + 1$ is a factor | M1 A1 cso | 2 | p(-1) attempted; not long division correctly shown = 0 plus statement |
| (c)(i) | Quadratic factor $(x^2 - 3x + 3)$ | M1 | | $b = -3$ or $c = 3$ by inspection or full long division attempt or comparing coefficients |
| | $(p(x) =) (x + 1)(x^2 - 3x + 3)$ | A1 | 2 | must see correct product |
| (ii) | Discriminant of quadratic $b^2 - 4ac = (-3)^2 - 4 \times 3$ | M1 | | 'their' discriminant considered possibly within quadratic equation formula |
| | $b^2 - 4ac < 0 \Rightarrow$ no real roots from quadratic \Rightarrow only one real root | A1 cso | 2 | |
| | Total | | 8 | |
| 6(a) | $\int_{-1}^1 (x^3 - 2x^2 + 3) dx$ $= \left[\frac{x^4}{4} - \frac{2x^3}{3} + 3x \right]_{-1}^1$ $= \left(\frac{1}{4} - \frac{2}{3} + 3 \right) - \left(\frac{1}{4} + \frac{2}{3} - 3 \right)$ $= 4\frac{2}{3}$ | M1 A1 A1 B1 ✓ A1 cso | 5 | one term correct another term correct all correct (condone + c) 'their' $F(1) - F(-1)$ with $(-1)^3$ etc evaluated correctly but must have earned M1 $\frac{14}{3}, \frac{56}{12}$ etc but combined as single fraction |
| (b) | Area of $\Delta \left(= \frac{1}{2} \times 2 \times 2 \right)$ $= 2$ Shaded region has area $4\frac{2}{3} - 2$ $= 2\frac{2}{3}$ | B1 M1 A1 cso | 3 | PI \pm their (a) \pm their Δ area $\frac{8}{3}, \frac{32}{12}$ etc but combined as single fraction |
| | Total | | 8 | |

MPC1 (cont)

| Q | Solution | Marks | Total | Comments |
|-------------|---|-------|----------|---|
| 7(a) | $8 - 6x > 5 - 4x - 8$ | M1 | 2 | multiplying out correctly and $>$ sign used |
| | $11 > 2x$ | A1cso | | accept $5.5 > x$ OE |
| | $x < 5\frac{1}{2} \quad \left(\text{or } x < \frac{11}{2} \right)$ | | | |
| | (b) $2x^2 + 5x - 12 \geq 0$ | | | |
| | $(x + 4)(2x - 3)$ | M1 | | correct factors |
| | Critical values are -4 and $\frac{3}{2}$ | A1 | | (or roots unsimplified) $\frac{-5 \pm \sqrt{121}}{4}$ both CVs correct; condone $\frac{6}{4}, -\frac{16}{4}$ etc here but must be single fractions |
| |  | M1 | | sketch or sign diagram including values |
| | $x \leq -4, \quad x \geq \frac{3}{2}$ | A1 | 4 | fractions must be simplified condone use of OR but not AND |
| | take their final line as their answer | | | |
| | Total | | 6 | |

MPC1 (cont)

| Q | Solution | Marks | Total | Comments |
|---------------|---|---------------------------|-----------|---|
| 8(a) | $(x-3)^2 + (y+8)^2 = 100$ | B1 B1 | 2 | accept $(y-8)^2$ condone $\text{RHS} = 10^2$ or $k = 10^2$ |
| (b) | $y=0 \Rightarrow \text{'their'}(x-a)^2 + b^2 = k$ $(x-3)^2 = 36$ or $x^2 - 6x - 27 (=0)$ (PI) $\Rightarrow x = -3, 9$ | M1 A1 A1 | 3 | Alternative  $(d^2 =) 10^2 - 8^2$ M1 $d^2 = 36$ A1 or $d = 6$ $\Rightarrow x = -3, 9$ A1 |
| (c) | Line CA has gradient $-\frac{2}{5}$ CA has equation $(y+8) = -\frac{2}{5}(x-3)$ $2x + 5y + 34 = 0$ | M1 A1 A1cso | 3 | any form of correct equation eg $y = -\frac{2}{5}x + c$, $c = -\frac{34}{5}$ integer coefficients - all terms on 1 side |
| (d)(i) | their $(x-3)^2 + (2x+1+8)^2$ or $x^2 + (2x+1)^2 - 6x + 16(2x+1)$ (+73) $x^2 - 6x + 9 + 4x^2 + 36x + 81 = 100$ or $x^2 + 4x^2 + 4x + 1 - 6x + 32x + 16 + 73 = 100$ $\Rightarrow 5x^2 + 30x - 10 = 0$ $\Rightarrow x^2 + 6x - 2 = 0$ | M1 A1 A1cso | 3 | substituting $y = 2x + 1$ correctly into LHS of "their" circle equation and attempt to expand in terms of x only any correct equation (with brackets expanded) must see this line or equivalent AG; all algebra must be correct |
| (ii) | $(x+3)^2 = 11$ $x = -3 \pm \sqrt{11}$ | M1 A1cso | 2 | or correct use of formula must get as far as $x = \frac{-6 \pm \sqrt{44}}{2}$ exactly this |
| | Total | | 13 | |
| | TOTAL | | 75 | |