



## General Certificate of Education

# Mathematics 6360 Statistics 6380

*MS/SS1A Statistics 1A*

## Mark Scheme

*2006 examination – January series*

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation meeting attended by all examiners and is the scheme which was used by them in this examination. The standardisation meeting ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for the standardisation meeting each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed at the meeting and legislated for. If, after this meeting, examiners encounter unusual answers which have not been discussed at the meeting they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

## Key To Mark Scheme And Abbreviations Used In Marking

M	mark is for method		
m or dM	mark is dependent on one or more M marks and is for method		
A	mark is dependent on M or m marks and is for accuracy		
B	mark is independent of M or m marks and is for method and accuracy		
E	mark is for explanation		
✓ or ft or F	follow through from previous incorrect result	MC	mis-copy
CAO	correct answer only	MR	mis-read
CSO	correct solution only	RA	required accuracy
AWFW	anything which falls within	FW	further work
AWRT	anything which rounds to	ISW	ignore subsequent work
ACF	any correct form	FIW	from incorrect work
AG	answer given	BOD	given benefit of doubt
SC	special case	WR	work replaced by candidate
OE	or equivalent	FB	formulae book
A2,1	2 or 1 (or 0) accuracy marks	NOS	not on scheme
–x EE	deduct x marks for each error	G	graph
NMS	no method shown	c	candidate
PI	possibly implied	sf	significant figure(s)
SCA	substantially correct approach	dp	decimal place(s)

### No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded. However, there are situations in some units where part marks would be appropriate, particularly when similar techniques are involved. Your Principal Examiner will alert you to these and details will be provided on the mark scheme.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

**Otherwise we require evidence of a correct method for any marks to be awarded.**

## MS/SS1A

Q	Solution	Marks	Total	Comments
1(a)	Gradient, $b = 0.886$ to $0.887$ $b = 0.88$ to $0.89$	B2 (B1)		AWFW AWFW
	Intercept, $a = 2.31$ to $2.33$ $a = 2.3$	B2 (B1)		AWFW AWRT
	<b>Alternatively</b> Attempt at $\Sigma x$ $\Sigma x^2$ $\Sigma y$ $\Sigma xy$			72, 624, 87, 720
	<b>or</b> Attempt at $S_{xx}$ $S_{xy}$	(M1)		105.6, 93.6
	Attempt at a correct formula for $b$ $b = 0.886$ to $0.887$ $a = 2.31$ to $2.33$	(m1) (A1) (A1)		AWFW AWFW
	Accept $a$ & $b$ interchanged only if $y = ax + b$ stated or subsequently used correctly in either (b) or (c)		4	
	(b) $a$ : average <b>waiting time</b> of 2.32 minutes (139 seconds) when entering <b>empty</b> <b>restaurant</b>	B1		OE; accept minimum waiting time
	$b$ : average <b>increase in waiting time</b> of 0.886 minutes (53 seconds) <b>for each</b> <b>customer in restaurant</b> on entry	B1	2	OE
	(c) Use of $y = a + 5b$ or $y = a + 25b$	M1		
	(i) For $x = 5$ $y = 6.6$ to $6.8$			
(d)(i)	(ii) For $x = 25$ $y = 24.3$ to $24.6$	A1	2	Both; AFWW
	Reliable as interpolation and small residuals	B1 B1		Within range OE OE
	<b>or</b> Reliable as interpolation but large %age residuals so inconclusive	(B1) (B1)		
	<b>or</b> Large %age residuals so unreliable	(B1)		
	(ii) Unreliable as extrapolation	B1	3	Outside range OE
	<b>Total</b>		<b>11</b>	

## MS/SS1A (cont)

Q	Solution	Marks	Total	Comments
<b>2(a)</b>	$P(X) = 0.3 \quad P(Y) = 0.4 \quad P(Z) = 0.2$			
<b>(i)</b>	$P(X \cap Y \cap Z) = 0.3 \times 0.4 \times 0.2$ $= 0.024$	M1	1	
<b>(ii)</b>	$P(X' \cap Y' \cap Z') = 0.7 \times 0.6 \times 0.8$ $= 0.336$	M1 A1	2	At least 2 correct terms CAO
<b>(iii)</b>	$P(X' \cap Y' \cap Z) + P(X' \cap Y \cap Z') +$ $P(X \cap Y' \cap Z') =$  $(0.7 \times 0.6 \times 0.2) + (0.7 \times 0.4 \times 0.8) +$ $(0.3 \times 0.6 \times 0.8) =$  $0.084 + 0.224 + 0.144 = 0.452$	M1  A1  A1	  3	At least two different combinations of three events  At least one correct numerical expression  CAO
<b>(b)</b>	$P(W   Z) = 0.9 \quad P(W   Z') = 0.25$			
<b>(i)</b>	$P(Z \cap W) = 0.2 \times 0.9$ $= 0.18$	M1 A1	2	Correct numerical expression CAO
<b>(ii)</b>	$P((Z \cap W') \cup (Z' \cap W))$ <b>or</b> $1 - [P((Z \cap W) \cup (Z' \cap W'))]$  $= 0.2 \times (1 - 0.9)$ $+$ $(1 - 0.2) \times 0.25$  $= 0.02 + 0.20$ $= 0.22$	M1  M1  A1	  3	$0.2 \times 0.9$ or (b)(i) $(1 - 0.2) \times (1 - 0.25)$  Cannot score an M1 in both methods  $1 - (0.18 + 0.60)$ CAO
	<b>Total</b>		<b>11</b>	


## MS/SS1A (cont)

Q	Solution	Marks	Total	Comments
3(a)	Mean = $\frac{286.5}{50} = 5.73$	B1	2	CAO
	Standard deviation = $\sqrt{\frac{45.16}{49 \text{ or } 50}} =$  0.95 to 0.961	B1		AWFW
	(b) 99% $\Rightarrow z = 2.57$ to 2.58	B1	4	AWFW 2.5758
	CI for $\mu$ is $\bar{x} \pm z \times \frac{(\sigma \text{ or } s)}{\sqrt{n}}$	M1		Use of Must have $(\div \sqrt{n})$ with $n > 1$
	Thus $5.73 \pm 2.5758 \times \frac{(0.95 \text{ to } 0.961)}{\sqrt{50}}$	A1✓		✓ on $z$ and $s^2 > 0$ but not on $\bar{x}$ Accept only 50 or 49 for $n$
	$5.73 \pm (0.34 \text{ to } 0.36)$	↑		Dependent
	(5.37 to 5.39, 6.07 to 6.09)	A1		AWFW
	(c) CI excludes both values of 5 and 6½ so Neither claim appears valid	B1✓ ↑ B1✓	2	✓ on (b); OE Dependent ✓ on (b); OE
	or			
	CI excludes 5 so claim not valid and CI excludes 6½ so claim not valid	(B1✓) (B1✓)		✓ on (b); OE ✓ on (b); OE
Total			8	

**MS/SS1A (cont)**

<b>Q</b>	<b>Solution</b>	<b>Marks</b>	<b>Total</b>	<b>Comments</b>
<b>4(a)</b>	Scatter Diagram <b>or</b> <b>or</b>	B2 (B1) (B1)	2	4 labelled points plotted 3 labelled points plotted 4 unlabelled points plotted
<b>(b)(i)</b>	Positive/linear correlation/relationship except for <b>two unusual</b> values/results	B1 B1	2	OE OE
<b>(ii)</b>	0.462	B1	1	CAO; accept 3 <sup>rd</sup> /final/last value
<b>(c)</b>	C and D C is likely freestyle champion D is likely backstroke champion <b>or</b> C is likely freestyle champion D is likely backstroke champion	B1 B1 (B1) (B1)	2	CAO Style identified
<b>(d)(i)</b>	$r = 0.912$ to $0.913$ <b>or</b> $r = 0.91$ to $0.92$ or $0.46$ to $0.47$ <b>or</b> $r = 0.9$ <b>Alternatively</b> Attempt at $\Sigma x \quad \Sigma x^2$ $\Sigma y \quad \Sigma y^2$ $\Sigma xy$ <b>or</b> Attempt at $S_{xx} \quad S_{yy} \quad S_{xy}$ Attempt at a correct formula for $r$ $r = 0.912$ to $0.913$	B3 (B2) (B1) (M1) (m1) (A1)	3	AWFW AWFW AWRT 270.4, 9188.46 301.6, 11437.84 10246.53 48.94, 67.52, 52.45
<b>(ii)</b>	Boys are faster/slower at both strokes <b>or</b> Boys are equally good at both strokes	B1	1	OE;do not accept freestyle times are proportional to backstroke times
<b>Total</b>			<b>11</b>	

## Question 4 (a)



(a) Scatter Diagram

4 labelled points plotted

B2

3 labelled points plotted

(B1)

4 unlabelled points plotted

(B1)

Total = 2

**MS/SS1A (cont)**

<b>Q</b>	<b>Solution</b>	<b>Marks</b>	<b>Total</b>	<b>Comments</b>
<b>5(a)(i)</b>	Time, $X \sim N(55, 8^2)$			
	$P(X < 60) = P\left(Z < \frac{60 - 55}{8}\right)$	M1		Standardising (59.5, 60 or 60.5) with 55 and ( $\sqrt{8}$ , 8 or $8^2$ ) and/or ( $55 - x$ )
	$= P(Z < 0.62 \text{ to } 0.63)$	A1		AWFW; ignore sign 0.625
	$= 0.732 \text{ to } 0.736$	A1	3	AWFW 0.73401
	<b>(ii)</b> $P(55 < X < 60) =$			
	$(i) - P(X < 55) = (i) - 0.5$	M1		Difference OE
	$= 0.232 \text{ to } 0.236$	A1	2	AWFW 0.23401
	<b>(b)</b> Time, $Y \sim N(\mu, 16^2)$			
	$0.95 \Rightarrow z = 1.64 \text{ to } 1.65$	B1		AWFW 1.6449
	$P(Y < 90) = P\left(Z < \frac{90 - \mu}{16}\right)$	M1		Standardising 90 using $\mu$ and 16
	Thus $\frac{90 - \mu}{16} = 1.6449$	m1		Equating z-term to z-value; Not using 0.95, 0.05 or $ 1 - z $
	Thus $\mu = 63.6 \text{ to } 63.8$	A1	4	AWFW
	<b>Total</b>		<b>9</b>	



## MS/SS1A (cont)

Q	Solution	Marks	Total	Comments
<b>6(a)(i)</b>	$B(20, p)$	M1	2	Use of in (a)
	$P(O \leq 10 \mid p = 0.40) = 0.872 \text{ to } 0.873$	A1		AWFW 0.8725
	<b>(ii)</b> $P(A = 3) = \binom{20}{3} (p)^3 (1-p)^{17}$	M1		Any value of $p$ with $0 < p < 1$
	$P(A = 3 \mid p = 0.28) =$		3	Fully correct expression
	$\binom{20}{3} (0.28)^3 (0.72)^{17}$	A1		
	$= 0.093 \text{ to } 0.095$	A1		
	<b>(iii)</b> $P(4 < B < 8 \mid p = 0.20) =$			AWFW 0.3383
		$P(B \leq 7 \text{ or } 8) = 0.9679 \text{ or } 0.9900$	M1	
		minus $P(B \leq 4 \text{ or } 3) = 0.6296 \text{ or } 0.4114$	M1	
		$= 0.338 \text{ to } 0.34$	A1	
		<b>or</b>		
<b>(b)</b>	Mean, $\mu = np = 500 \times 0.12 = 60$	B1	3	Or implied by a correct answer
	Variance, $\sigma^2 = np(1-p) = 60 \times 0.88 = 52.8$	B1		
			2	CAO
	<b>Total</b>		<b>10</b>	
	<b>TOTAL</b>		<b>60</b>	