

# **General Certificate of Education June 2010**

Mathematics

**MS03** 

**Statistics 3** 

Mark Scheme

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation meeting attended by all examiners and is the scheme which was used by them in this examination. The standardisation meeting ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for the standardisation meeting each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed at the meeting and legislated for. If, after this meeting, examiners encounter unusual answers which have not been discussed at the meeting they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available to download from the AQA Website: www.aqa.org.uk

Copyright © 2010 AQA and its licensors. All rights reserved.

#### **COPYRIGHT**

AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

## Key to mark scheme and abbreviations used in marking

M	mark is for method				
m or dM	mark is dependent on one or more M marks and is for method				
A	mark is dependent on M or m marks and is for accuracy				
В	mark is independent of M or m marks and is for method and accuracy				
E	mark is for explanation				
√or ft or F	follow through from previous				
	incorrect result	MC	mis-copy		
CAO	correct answer only	MR	mis-read		
CSO	correct solution only	RA	required accuracy		
AWFW	anything which falls within	FW	further work		
AWRT	anything which rounds to	ISW	ignore subsequent work		
ACF	any correct form	FIW	from incorrect work		
AG	answer given	BOD	given benefit of doubt		
SC	special case	WR	work replaced by candidate		
OE	or equivalent	FB	formulae book		
A2,1	2 or 1 (or 0) accuracy marks	NOS	not on scheme		
–x EE	deduct x marks for each error	G	graph		
NMS	no method shown	c	candidate		
PI	possibly implied	sf	significant figure(s)		
SCA	substantially correct approach	dp	decimal place(s)		

### No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded. However, there are situations in some units where part marks would be appropriate, particularly when similar techniques are involved. Your Principal Examiner will alert you to these and details will be provided on the mark scheme.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

Otherwise we require evidence of a correct method for any marks to be awarded.

## **MS03**

Q	Solution	Marks	Total	Comments
1	$H_0$ : $\rho = 0$	В1		Both
	$H_1: \rho \neq 0$	21		
	$SL \qquad \alpha = 0.05 (5\%)$			
	` /	D.1		AWRT (0.4044)
	$CV   r = (\pm) 0.404$	B1		$H_1: \rho > 0 \Rightarrow r = 0.3438$
	Calculated $r = 0.336$ < Tabulated $r$	M1		Comparison
	No evidence, at 5% level, of a			F on CV
	correlation between stem length and cup	A1F	4	At 5% level, accept hypothesis of no
	diameter of matsutake mushrooms	1111	•	correlation
	Total		4	
2(a)	99% $\Rightarrow z = 2.57 \text{ to } 2.58$	B1		AWFW (2.5758)
		3.64		
	CI for $\mu_R - \mu_D$ is	M1		Form
	CI for $\mu_R - \mu_D$ is $(\overline{x}_R - \overline{x}_D) \pm z \times \sqrt{\frac{s_R^2 + s_D^2}{n_D}}$			Allow $\left(\frac{ns^2}{n-1}\right)$ or $(n-1)$
	$(x_R - x_D) \pm z \times \sqrt{\frac{n}{n_B} + \frac{z}{n_D}}$			
	, K D	A1		Correct expression
	[ <del>-2</del> -2			
	ie $(225-219) \pm 2.5758 \sqrt{\frac{5^2}{50} + \frac{8^2}{75}}$	A1F		Or equivalent F on z only
	V 50 75	7111		r on 2 only
	ie $6 \pm 3$ or $(3, 9)$	A1	5	CAO/AWRT or AWRT
	Note:	111		
	Use of pooled $s^2 = 5961/123 = 48.46341 \Rightarrow$			
	$6 \pm 3.3 \Rightarrow \text{max of B1 M1 A0 A1F A0 (3)}$			
<b>a</b> >		Die		
(b)	CI does not include 0/zero	B1F		F on (a)
	Evidence of a <b>difference</b> in mean weights	B1F		F on (a)
		dep	2	Dependent on CI but not on 0/zero
(c)	Price, size, quality, taste, presentation,	B1	1	Any sensible reason
	organic, marketing, stall position, etc			,
	Total		8	

MS03 (cont)				
Q	Solution	Marks	Total	Comments
3	$H_0$ : $\lambda_T = \lambda_S$ $H_1$ : $\lambda_T > \lambda_S$	B1		Both
	SL $\alpha = 0.02 (2\%)$	D.1		AWEW (2.0527)
	CV $z = 2.05 \text{ to } 2.06$	B1		AWFW (2.0537)
	or $H_1 \lambda_T \neq \lambda_s \implies z = 2.32 \text{ to } 2.33$	(B1)		AWFW (2.3263)
	$\overline{s} = \frac{940}{40} = 23.5$ $\overline{t} = \frac{1560}{60} = 26$	B1		Both CAO; may be implied
	Pooled value, $\bar{p} = \frac{2500}{100} = 25$	B1		CAO
	$z = \frac{ 23.5 - 26 }{\sqrt{25\left(\frac{1}{40} + \frac{1}{60}\right)}} \text{ or } z = \frac{ 23.5 - 26 }{\sqrt{\left(\frac{23.5}{40} + \frac{26}{60}\right)}}$	M1		
	z = 2.44 to 2.45 or $z = 2.47$ to 2.48	A1		Either AWFW (2.449 or 2.474)
	<b>Evidence</b> , at 2% level, <b>to agree</b> with Tina's claim	A1F	7	F on CV and z-value
	Total		7	

IS03 (cont) Q	Solution	Marks	Total	Comments
4(a)	A+(0.90) 0.09	B1		S and NS with Ps or %s
( )	- S(0.10)——A-(0.02) 0.002			
		B1		$2 \times (A+ \text{ and } A-) \text{ with Ps or } \%s$
	B(0.08)—— +(0.98) 0.00784	D.1		
		B1		$2 \times (B)$ with Ps or %s
	A+(0.01)	B1		$2 \times (B+ \text{ and } B-) \text{ with Ps or } \%s$
	NS(0.90)A-(0.80) 0.72	D1		2 × (B+ and B=) with 13 of 703
	B(0.19)—— $+(0.01)$ 0.00171	(B2,1)	4	Basic shape with labels, but without Ps
	-(0.99) 0.16929			or %s
	Note:			
	The following BF and AF marks are dependent on an essentially correctly-			
	shaped tree diagram			
	end and and			
(b)(i)				
(A)	P(S  and  -) = 0.002 + 0.00016 = 0.00216	B1F		F on (a); otherwise CAO
( <b>D</b> )	P(NS  and  +) = 0.009 + 0.00171 = 0.01071	B1F	2	F on (a); otherwise AWRT 0.0107
<b>(B)</b>	F(NS  and  +) = 0.009 + 0.00171 = 0.01071	БІГ	2	r on (a), otherwise AWK1 0.0107
(ii)	$E(N) = 10000 \times [(A) + (B)]$	M1		Or equivalent
, ,	200			•
	= $128.6$ to $128.7 \Rightarrow 130$	A1F	2	CAO
(a)(i)	D(C 1.)			
(c)(i)	$P(S \mid +) = \frac{P(S \text{ and } +)}{P(+)} =$	M1		Used
	· /	M1		
	$\frac{0.09 + 0.00784}{0.00784} = \frac{0.09784}{0.00784}$	A1F		F on (a)
	0.09 + 0.00784 + 0.009 + 0.00171 $0.10855$	7111		Otherwise correct
	0.001 / 0.002	4.1		A NADE (0.0012.4)
	= 0.901 to 0.902	A1		AWRT (0.90134)
(ii)	P(NS and -)			
(11)	$P(NS \mid -) = \frac{P(NS \text{ and } -)}{P(NS \mid -)} =$	(M1)		Used; only if not scored in (i)
	P(-)			
	$\frac{0.72 + 0.16929}{0.00000000000000000000000000000000000$	A1F		F on (a) and/or denominator (c)(i)
	0.002 + 0.00016 + 0.72 + 0.16929 $0.89145$			Otherwise correct
	= 0.997 to 0.998	A1	5	AWFW (0.99758)
	Special cases:	111		(0.77730)
	Only numerators correct $\Rightarrow$ (M1) B1 B1			
	Only denominators correct $\Rightarrow$ (M1) B1 B1			
	Total		13	

MS03 (cont)		N	TD 4 1		
Q <b>5</b> (a)	Solution 5.77 - 2 × 250 + 2 × 210 - 1120	Marks	Total	CAO	
5(a)	$E(T) = 2 \times 350 + 2 \times 210 = 1120$	B1		CAO	
	$Cov(W, H) = \sqrt{5 \times 4} \times 0.75 =$	M1		Used; may be implied	
	3.34 to 3.36	<b>A</b> 1		AWFW	(3.3541)
	Var $(T) = (2^2 \times 5) + (2^2 \times 4) + (2 \times 2 \times 2 \times 3.3541)$	M1		Used Ignore 3rd expression	
	= 20 + 16 + 26.8328 = 62.7 to 62.9	A1	5	AWFW	(62.8328)
(b)	$L = T_1 + T_2 + T_3 + T_4$				
	Mean of $L = 4480$	B1F		CAO; F on $E(T)$	
	Variance of $L = 4 \times Var(T)$	M1			
	= 250.8 to 251.6				(251.3312)
	SD of $L = 15.8$ to 15.9	A1F		Either AWFW; F on Var (T)	(15.8534)
	$P(L < 4500) = P\left(Z < \frac{4500 - 4480}{\sqrt{251.3312}}\right)$	M1		Standardising 4500 using C's mean and SD	
	= P(Z < 1.25  to  1.27)				
	= 0.894 to 0.898	A1		AWFW	(0.89645)
	Alternative Solution: Use of $\overline{T}$ rather than $L$				
	Mean of $\overline{T} = 1120$	(B1F)		CAO; F on $E(T)$	
	Variance of $\overline{T} = Var(T) \div 4$	(M1)			
	= 15.6 to 15.8	(A1F)		Either AWFW; F on Var(T)	(15.7082)
	SD of $\bar{T} = 3.95$ to 3.97	(AII <sup>*</sup> )		Limei Awrw, i on val(1)	(3.9634)
	$P(\overline{T} < 1125) = P\left(Z < \frac{1125 - 1120}{\sqrt{15.7082}}\right)$	(M1)		Standardising 1125 using C's mean and SD	
	= P(Z < 1.25  to  1.27)				
	= 0.894 to 0.898	(A1)	5	AWFW	(0.89645)
	Total		10		

MS03 (cont)				<del>,</del>
Q	Solution	Marks	Total	Comments
6(a)(i)	$\hat{p} = \frac{28}{175} = 0.16$	B1		CAO; or equivalent
	$95\% \Rightarrow z = 1.96$	B1		AWRT
	Approximate CI for $p$ is $\hat{p} \pm z \times \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$	M1		Used
	ie $0.16 \pm 1.96 \sqrt{\frac{0.16 \times 0.84}{175}}$	A1F		Or equivalent F on $\hat{p}$ and z
	ie $0.16 \pm 0.054$ or $(0.106, 0.214)$	<b>A</b> 1	5	CAO/AWRT or AWRT (0.0543)
(ii)	<b>CI does</b> include <b>0.2</b> (20%)	B1F		F on (i)
	No evidence to support councils' claim	B1F dep	2	F on (i) Dependent on CI and on 0.2
(b)(i)	$H_0$ : $p = 0.40 (40\%)$ $H_1$ : $p < 0.40$	B1		Both
	Using B (50, 0.4) (40%)	M1		May be implied
	$P(X \le 16) = 0.156$	A1		AWRT (0.1561)
	Calculated probability > 0.10 (10%)	M1		Comparison
	No evidence, at 10% level, to support council's claim  Special Case: Normal approximation $z = -1.15(47) \text{ B1}$ CV = -1.28(16) B1  Conclusion B1F Max of 4 marks	A1F	5	F on probability v 0.10 or 0.05 At 10% level, <b>accept</b> (at least) <b>40%</b> Allow B1 for hypotheses p = 0.123 to 0.125 v 0.10 B1 B1 F on z and CV
(ii)	Require $P(X \le x) \le 0.10$	M1		May be implied Ignore any reasoning if '15' stated
	$\Rightarrow$ CV = 15 (CR $\leq$ 15)	A1	2	CAO; or equivalent
(iii)	$P(Type II error) = P(accept H_0   H_0 false)$	B1		Stated or used; or equivalent
	$= P(X > CV \text{ or } X \ge CV)$	M1		Attempt at a probability > or ≥ C's CV from (ii)
	= 1 - (0.8369  or  0.7481)	M1		Ignore '1 -'
	= 0.163	A1	4	AWRT
	Total		18	

MS03 (cont)				
Q	Solution	Marks	Total	Comments
7	$X \sim \text{Po}(\lambda)$			
(a)(i)	$E(X) = \sum_{x=0}^{\infty} x \times \frac{e^{-\lambda} \lambda^x}{x!} =$	M1		Used; ignore limits until A1
	$\lambda e^{-\lambda} \times \sum_{x=1}^{\infty} \frac{\lambda^{x-1}}{(x-1)!} =$	M1		Factor of at least $\lambda$ Division of $x$ ! by $x$
	$\lambda e^{-\lambda} \times e^{\lambda} = \lambda$	A1	3	AG; fully correct solution
(ii)	$Var(X) = E(X^{2}) - [E(X)]^{2}$ = E[X(X-1)] + E(X) - [E(X)]^{2}	M1		Used (Other derivations are possible)
	$= \lambda^2 + \lambda - \lambda^2 = \lambda$	A1	2	CAO
(b)(i)	$P(X = m) \ge P(X = m - 1)$ and $P(X = m) \ge P(X = m + 1) \Rightarrow$			
	$\frac{\mathrm{e}^{-\lambda}\lambda^m}{m!} \ge \frac{\mathrm{e}^{-\lambda}\lambda^{m-1}}{(m-1)!} \text{ and } \frac{\mathrm{e}^{-\lambda}\lambda^m}{m!} \ge \frac{\mathrm{e}^{-\lambda}\lambda^{m+1}}{(m+1)!}$	M1		Use of $Po(\lambda)$ for $x = m$
	m! $(m-1)!$ $m!$ $(m+1)!$	M1		Either inequality (accept = sign)
	$m \le \lambda$ and $m \ge \lambda - 1$	A1	3	AG; fully correct solution
(ii)	Given $\lambda = 4.9 \implies m = 4$	B1		CAO
	$P(X=4) = \frac{e^{-4.9} 4.9^4}{4!} = 0.178 \text{ to } 0.179$	B1	2	AWFW (0.178867)
(c)	Given SD $(Y) = 15.5 \implies \lambda = \text{Var}(Y) = 15.5^2 = 240.25$	B1		Either CAO
	Mode, $d = 240$	B1F		F on $\lambda$ providing an integer value
	$P(Y_P \ge d) = P(Y_N > d - 0.5) =$	В1		Accept use of 'd'
	$P\bigg(Z > \frac{239.5 - 240.25}{15.5}\bigg) =$	M1		Standardising $(d-0.5, d \text{ or } d+0.5)$ with $15.5^2$ and $15.5$ ; do <b>not</b> accept use of 'd'
	P(Z > -0.05) = 0.515  to  0.52	A1	5	AWFW (0.5193)
	Total		15	
	TOTAL		75	
	P(Z > -0.05) = 0.515  to  0.52		15	$15.5^2$ and $15.5$ ; do <b>not</b> accept use of 'd'