

**Physics (B) Physics in Context**  
**Unit 1 Harmony and Structure in the Universe**

**PHYB1**


**Data and Formulae Booklet**

**FUNDAMENTAL CONSTANTS AND  
OTHER NUMERICAL DATA**

| Quantity                     | Symbol       | Value                          | Units                             |
|------------------------------|--------------|--------------------------------|-----------------------------------|
| speed of light in vacuo      | $c$          | $3.00 \times 10^8$             | $\text{m s}^{-1}$                 |
| Planck constant              | $h$          | $6.63 \times 10^{-34}$         | J s                               |
| gravitational constant       | $G$          | $6.67 \times 10^{-11}$         | $\text{N m}^2 \text{kg}^{-2}$     |
| gravitational field strength | $g$          | 9.81                           | $\text{N kg}^{-1}$                |
| acceleration due to gravity  | $g$          | 9.81                           | $\text{m s}^{-2}$                 |
| electron rest mass           | $m_e$        | $9.11 \times 10^{-31}$         | kg                                |
|                              | $m_e$        | $5.5 \times 10^{-4} \text{ u}$ |                                   |
| electron charge              | $e$          | $-1.60 \times 10^{-19}$        | C                                 |
| proton rest mass             | $m_p$        | $1.67(3) \times 10^{-27}$      | kg                                |
|                              | $m_p$        | 1.00728 u                      |                                   |
| neutron rest mass            | $m_n$        | $1.67(5) \times 10^{-27}$      | kg                                |
|                              | $m_n$        | 1.00867 u                      |                                   |
| permittivity of free space   | $\epsilon_0$ | $8.85 \times 10^{-12}$         | $\text{F m}^{-1}$                 |
| molar gas constant           | $R$          | 8.31                           | $\text{J K}^{-1} \text{mol}^{-1}$ |
| Boltzmann constant           | $k$          | $1.38 \times 10^{-23}$         | $\text{J K}^{-1}$                 |
| Avogadro constant            | $N_A$        | $6.02 \times 10^{23}$          | $\text{mol}^{-1}$                 |
| Wien constant                | $\alpha$     | $2.90 \times 10^{-3}$          | m K                               |

**GEOMETRICAL  
EQUATIONS**

|                          |                             |
|--------------------------|-----------------------------|
| arc length               | $r\theta$                   |
| circumference of circle  | $2\pi r$                    |
| area of circle           | $\pi r^2$                   |
| surface area of sphere   | $4\pi r^2$                  |
| volume of sphere         | $\frac{4}{3}\pi r^3$        |
| surface area of cylinder | $2\pi rh$                   |
| volume of cylinder       | $\pi r^2 h$                 |
|                          | $\sin \theta = \frac{a}{c}$ |
|                          | $\cos \theta = \frac{b}{c}$ |
|                          | $\tan \theta = \frac{a}{b}$ |
|                          | $c^2 = a^2 + b^2$           |



**Unit Conversions**

|                        |                                    |
|------------------------|------------------------------------|
| 1 atomic mass unit (u) | $1.661 \times 10^{-27} \text{ kg}$ |
| 1 year (y)             | $3.15 \times 10^7 \text{ s}$       |
| 1 parsec (pc)          | $3.08 \times 10^{16} \text{ m}$    |
| 1 parsec               | 3.26 ly                            |
| 1 light year (ly)      | $9.46 \times 10^{15} \text{ m}$    |

**Particle Properties**

**Properties of quarks** antiquarks have opposite signs

| type     | charge          | Baryon number  | strangeness |
|----------|-----------------|----------------|-------------|
| <b>u</b> | $+\frac{2}{3}e$ | $+\frac{1}{3}$ | 0           |
| <b>d</b> | $-\frac{1}{3}e$ | $+\frac{1}{3}$ | 0           |
| <b>s</b> | $-\frac{1}{3}e$ | $+\frac{1}{3}$ | -1          |

**Properties of Leptons**

|                                                                                    | Lepton Number |
|------------------------------------------------------------------------------------|---------------|
| particles:<br>$e^-, \nu_e; \mu^-, \nu_\mu; \tau^-, \nu_\tau$                       | +1            |
| antiparticles:<br>$e^+, \bar{\nu}_e; \mu^+, \bar{\nu}_\mu; \tau^+, \bar{\nu}_\tau$ | -1            |

## AS FORMULAE

| Waves                                                              |                                                                                               | Quantum Physics and Astrophysics          |                                                                       |
|--------------------------------------------------------------------|-----------------------------------------------------------------------------------------------|-------------------------------------------|-----------------------------------------------------------------------|
| wave speed                                                         | $c = f\lambda$                                                                                | photon energy                             | $E = hf$                                                              |
| period                                                             | $T = \frac{1}{f}$                                                                             | Einstein equation                         | $hf = \phi + E_{k(\max)}$                                             |
| intensity                                                          | $I = \frac{P}{A}$                                                                             | line spectrum equation                    | $hf = E_1 - E_2$                                                      |
| stretched string frequency                                         | $f = \frac{1}{2L} \sqrt{\frac{T}{\mu}}$                                                       | de Broglie wavelength                     | $\lambda = \frac{h}{p} = \frac{h}{mv}$                                |
| beat frequency                                                     | $f = f_1 - f_2$                                                                               | Doppler shift for $v \ll c$               | $\frac{\Delta f}{f} = -\frac{\Delta \lambda}{\lambda} = \frac{v}{c}$  |
| fringe spacing                                                     | $w = \frac{\lambda D}{s}$                                                                     | Wien's law                                | $\lambda_{\max} T = 0.0029 \text{ m K}$                               |
| diffraction grating                                                | $n\lambda = d \sin \theta$                                                                    | Hubble law                                | $v = Hd$                                                              |
| half beam width                                                    | $\sin \theta = \frac{\lambda}{a}$                                                             | intensity for a point source              | $I = \frac{P}{4\pi r^2}$                                              |
| refractive index of a substance (s)                                | $n = \frac{c}{c_s}$                                                                           | <b>Electricity</b>                        |                                                                       |
| for two different substances of refractive indices $n_1$ and $n_2$ | $n_1 \sin i_1 = n_2 \sin i_2$                                                                 | current                                   | $I = \frac{\Delta Q}{\Delta t}$                                       |
| critical angle                                                     | $\sin \theta_c = \frac{n_2}{n_1} \text{ for } n_1 > n_2$                                      | electromotive force (emf)                 | $\varepsilon = \frac{E}{Q}$                                           |
| <b>Mechanics</b>                                                   |                                                                                               |                                           | $\varepsilon = I(R + r)$                                              |
| speed or velocity                                                  | $v = \frac{\Delta s}{\Delta t}$                                                               | resistance                                | $R = \frac{V}{I}$                                                     |
| acceleration                                                       | $a = \frac{\Delta v}{\Delta t}$                                                               | resistors in series                       | $R = R_1 + R_2 + R_3 + \dots$                                         |
| equations of motion                                                | $v = u + at$<br>$s = \frac{(u + v)}{2} t$<br>$v^2 = u^2 + 2as$<br>$s = ut + \frac{1}{2} at^2$ | resistors in parallel                     | $\frac{1}{R} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3} + \dots$ |
| force                                                              | $F = ma$                                                                                      | resistivity                               | $\rho = \frac{RA}{L}$                                                 |
| change in potential energy                                         | $\Delta E_p = mg\Delta h$                                                                     | power                                     | $P = VI = I^2 R = \frac{V^2}{R}$                                      |
| kinetic energy                                                     | $E_k = \frac{1}{2} mv^2$                                                                      | potential divider formula                 | $V_o = \left( \frac{R_1}{R_1 + R_2} \right) \times V_i$               |
| momentum                                                           | $p = mv$                                                                                      | energy                                    | $E = VI t$                                                            |
| impulse                                                            | $F\Delta t = \Delta(mv)$                                                                      | efficiency                                | $\frac{\text{useful output power}}{\text{input power}}$               |
| spring stiffness                                                   | $k = \frac{F}{\Delta L}$                                                                      | <b>Energy production and transmission</b> |                                                                       |
| energy stored for $F \propto L$                                    | $E = \frac{1}{2} F\Delta L$                                                                   | rate of heat transfer by conduction       | $= UA \Delta \theta$                                                  |
| work done                                                          | $W = Fs \cos \theta$                                                                          | maximum power for a wind turbine          | $= \frac{1}{2} \pi r^2 \rho v^3$                                      |
| power                                                              | $P = \frac{\Delta W}{\Delta t} = Fv$                                                          |                                           |                                                                       |
| density                                                            | $\rho = \frac{m}{V}$                                                                          |                                           |                                                                       |