



**General Certificate of Education (A-level)  
January 2012**

**Mathematics**

**MFP2**

**(Specification 6360)**

**Further Pure 2**

**Final**

***Mark Scheme***

---

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all examiners participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for standardisation each examiner analyses a number of students' scripts: alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, examiners encounter unusual answers which have not been raised they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available from: [aqa.org.uk](http://aqa.org.uk)

Copyright © 2012 AQA and its licensors. All rights reserved.

**Copyright**

AQA retains the copyright on all its publications. However, registered schools/colleges for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

## Key to mark scheme abbreviations

M	mark is for method
m or dM	mark is dependent on one or more M marks and is for method
A	mark is dependent on M or m marks and is for accuracy
B	mark is independent of M or m marks and is for method and accuracy
E	mark is for explanation
✓ or ft or F	follow through from previous incorrect result
CAO	correct answer only
CSO	correct solution only
AWFW	anything which falls within
AWRT	anything which rounds to
ACF	any correct form
AG	answer given
SC	special case
OE	or equivalent
A2,1	2 or 1 (or 0) accuracy marks
-x EE	deduct x marks for each error
NMS	no method shown
PI	possibly implied
SCA	substantially correct approach
c	candidate
sf	significant figure(s)
dp	decimal place(s)

## No Method Shown


Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded.


Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

**Otherwise we require evidence of a correct method for any marks to be awarded.**

Q	Solution	Marks	Total	Comments
1(a)	 <p data-bbox="240 674 448 705">Sketch <math>y = \sinh x</math></p> <p data-bbox="240 748 464 779">Sketch <math>y = \operatorname{sech} x</math>:</p> <p data-bbox="240 790 703 822">Symmetry about <math>x=0</math> with max point</p> <p data-bbox="240 826 443 857">Asymptote <math>y=0</math></p> <p data-bbox="240 862 603 893">Point <math>(0, 1)</math> marked or implied</p>	B1		gradient $> 0$ at $(0, 0)$ ; no asymptotes
(b)	$\sinh x = \frac{1}{\cosh x}$ $\sinh 2x = 2$ <p data-bbox="240 1048 347 1079">Use of <math>\ln</math></p> $x = \frac{1}{2} \ln(2 + \sqrt{5})$ <p data-bbox="240 1162 272 1193"><b>or</b></p> $\frac{1}{2}(e^{2x} - e^{-2x}) = 2 \quad \text{OE}$ $e^{4x} - 4e^{2x} - 1 = 0$ <p data-bbox="240 1308 512 1339">Correct use of formula</p> <p data-bbox="240 1344 320 1375">Result</p>	M1 M1 m1 A1 (M1) (M1) (m1) (A1)	4 4 4 (4)	must not cross $x$ -axis  use of double angle formula dependent on previous M2  incorrect $\sinh x$ , $\cosh x$ M0 (no marks) ie multiply by $e^{2x}$ and rewrite
	<b>Total</b>		<b>8</b>	

Q	Solution	Marks	Total	Comments
2(a)	 <p>Half-line with gradient <math>&lt; 1</math></p>	B1	1	condone a short line, ie it stops at or inside circle
(b)(i)	Circle centre on $L$ , $x$ -coord 6 indicated touching $\text{Re } z = 0$ not at $(0, 0)$	B1 B1	2	not touching $\text{Re}$ axis
(ii)	<p><math>y</math>-coord of centre is <math>2\sqrt{3}</math> or <math>\frac{6}{\sqrt{3}}</math></p> <p><math>z_0 = 6 + 2\sqrt{3}i</math>, <math>k = 6</math></p>	B1 B1F, B1	3	OE; PI ft error in coords of centre
(iii)	<p>Point <math>z_1</math> shown</p> <p><math>\arg z_1 = -\frac{1}{6}</math></p>	B1 B1	2	PI
<b>Total</b>			<b>8</b>	
3(a)	$\frac{dy}{dx} = \frac{1}{2 \tanh x}$ $\times \text{sech}^2 x$ $= \frac{1}{2 \sinh x \cosh x}$ $= \frac{1}{\sinh 2x}$	B1 B1 M1 A1	4	for expressing in terms of $\sinh x$ and $\cosh x$ AG; PI by previous line
(b)	$\sqrt{1 + \left(\frac{dy}{dx}\right)^2} = \sqrt{1 + \frac{1}{\sinh^2 2x}}$ $= \sqrt{\frac{\cosh^2 2x}{\sinh^2 2x}}$ $= \frac{\cosh 2x}{\sinh 2x}$ <p>Integral is <math>\frac{1}{2} \ln \sinh 2x</math></p> $\sinh(2 \ln 4) = \frac{255}{32} \quad \sinh(2 \ln 2) = \frac{15}{8}$ $s = \frac{1}{2} \ln \left(\frac{17}{4}\right)$	M1 m1 A1 M1A1 B1B1 A1F	8	use of formula; accept $\sqrt{\quad}$ inserted at any stage relevant use of $\cosh^2 - \sinh^2 = 1$ OE M1 for $\ln \sinh$ PI ft error in $\frac{1}{2}$
<b>Total</b>			<b>12</b>	

Q	Solution	Marks	Total	Comments
4	Assume result true for $n = k$ Then $u_{k+1} = \frac{3}{4 - \left(\frac{3^{k+1} - 3}{3^{k+1} - 1}\right)}$ $= \frac{3(3^{k+1} - 1)}{4(3^{k+1} - 1) - (3^{k+1} - 3)}$ $4 \times 3^{k+1} - 3^{k+1} = 3^{k+2}$ $u_{k+1} = \frac{3^{k+2} - 3}{3^{k+2} - 1}$ $n = 1 \quad \frac{3^2 - 3}{3^2 - 1} = \frac{3}{4} = u_1$ Induction proof set out properly	M1 A1 A1 A1 B1 E1	6	clearly shown  must have earned previous 5 marks
<b>Total</b>			<b>6</b>	
5	Numerator = $e^{\frac{p\pi i}{8}}$ Denominator = $e^{\frac{-q\pi i}{12}}$ Fraction = $e^{\frac{p\pi i}{8} + \frac{q\pi i}{12}}$ $= e^{\frac{\pi i}{24}(3p+2q)}$ $i = e^{\frac{12\pi i}{24}}$ $3p + 2q = 12$ $p = 2, q = 3$ <p><b>Alternative 1</b></p> Numerator = $\cos \frac{p\pi}{8} + i \sin \frac{p\pi}{8}$ Denominator = $\cos \frac{-q\pi}{12} + i \sin \frac{-q\pi}{12}$ Fraction = $\left(\cos \frac{p\pi}{8} + i \sin \frac{p\pi}{8}\right) \left(\cos \frac{q\pi}{12} + i \sin \frac{q\pi}{12}\right)$ $= \cos \frac{\pi}{24}(3p + 2q) + i \sin \frac{\pi}{24}(3p + 2q)$ $= i \text{ if } \cos \frac{\pi}{24}(3p + 2q) = 0$ $\text{or } \sin \frac{\pi}{24}(3p + 2q) = 1$ $3p + 2q = 12$ $p = 2, q = 3$ <p><b>Alternative 2</b></p> LHS $\cos \frac{p\pi}{8} + i \sin \frac{p\pi}{8}$ RHS $i \cos \frac{q\pi}{12} + \sin \frac{q\pi}{12}$ $\cos \frac{p\pi}{8} = \sin \frac{q\pi}{12}$ or $\sin \frac{p\pi}{8} = \cos \frac{q\pi}{12}$ Introduction of $\frac{\pi}{2}$ $\frac{p\pi}{8} = \frac{\pi}{2} - \frac{q\pi}{12}$ $3p + 2q = 12$ $p = 2, q = 3$	B1 B1 M1 A1 m1 A1F A1 (B1) (B1) (M1) (A1) (m1) (A1F) (A1) (B1) (B1) (M1) (m1) (A1) (A1F) (A1)	7	allow for attempt to subtract powers  OE ft errors of sign or arithmetic slips CAO  needs more than just $\cos \frac{q\pi}{12} - \sin \frac{p\pi}{12}$  CAO  CAO (correct answers, insufficient working 3/7 only)
<b>Total</b>			<b>7</b>	

Q	Solution	Marks	Total	Comments
6(a)	$7 + 4x - 2x^2 = 9 - 2(x-1)^2$	M1A1	2	
(b)	Put $u = \sqrt{2}(x-1)$ $du = \sqrt{2} dx$ $I = \frac{1}{\sqrt{2}} \int \frac{du}{\sqrt{9-u^2}}$ $= \frac{1}{\sqrt{2}} \sin^{-1} \frac{u}{3}$ Change limits or replace $u$ $= \frac{\pi}{4\sqrt{2}}$ or $\frac{\pi\sqrt{2}}{8}$	M1 A1F A1F A1 m1 A1	6	allow $u = k(x-1)$ any $k$ ft error in (a); must have $u^2$ only, ie $\frac{1}{\sqrt{2}}$ outside integrand for $\sin^{-1} \frac{u}{p}$ provided $\sin^{-1}$
	<b>Alternative</b> – if integration is attempted without substitution: $\sin^{-1} \frac{1}{\sqrt{2}}$ $(x-1) \frac{\sqrt{2}}{3}$ Substitution of limits $\frac{\pi}{4\sqrt{2}}$	(M1) (A1F) (A1) (A1F) (m1) (A1)	(6)	CAO
<b>Total</b>			<b>8</b>	
7(a)	Use of $(\sum \alpha)^2 = \sum \alpha^2 + 2\sum \alpha\beta$	M1 A1	2	AG
(b)	$p = 0, q = 5 + 6i$	B1,B1	2	
(c)(i)	Substitute $3i$ for $z$ <b>or</b> use $3i\beta\gamma = -r$ $-27i + 15i - 18 + r = 0$ <b>or</b> $\beta\gamma = 5 + 6i + \alpha^2$ $r = 18 + 12i$	M1 A1 A1F	3	allow for $3i\beta\gamma = r$ any form one error
(ii)	Cubic is $(z-3i)(z^2 + 3iz - 4 + 6i)$ or use of $\beta\gamma$ and $\beta + \gamma$	M1A1	2	clearly shown
(iii)	$f(-2) = 0$ or equate imaginary parts $\beta = -2, \gamma = 2 - 3i$	M1 A1,A1F	3	correct answers no working and no check B1 only
<b>Total</b>			<b>12</b>	

Q	Solution	Marks	Total	Comments
8(a)	$1, e^{\frac{2\pi i}{5}}, e^{\frac{4\pi i}{5}}, e^{\frac{-2\pi i}{5}}, e^{\frac{-4\pi i}{5}}$	B1	1	accept $e^0$
(b)	$\frac{z^5 - 1}{z - 1} = z^4 + z^3 + z^2 + z + 1$ $= \left(z - e^{\frac{2\pi i}{5}}\right) \left(z - e^{\frac{4\pi i}{5}}\right) \left(z - e^{\frac{-2\pi i}{5}}\right) \left(z - e^{\frac{-4\pi i}{5}}\right)$	B1 M1A1	3	B0 if assumed accept if $e^{\frac{6\pi i}{5}}, e^{\frac{8\pi i}{5}}$ used here
(c)	Correct grouping of linear factors $e^{\frac{2\pi i}{5}} + e^{\frac{-2\pi i}{5}} = 2 \cos \frac{2\pi}{5}$ $\left(z^2 - 2 \cos \frac{2\pi}{5} z + 1\right) \left(z^2 - 2 \cos \frac{4\pi}{5} z + 1\right)$ $\div z^2$ to give answer	M1 A1 A1 A1	4	clearly shown AG
(d)	Substitute into LHS to give $w^2 + w - 1$ RHS $\left(w - 2 \cos \frac{2\pi}{5}\right) \left(w - 2 \cos \frac{4\pi}{5}\right)$ Solve $w^2 + w - 1 = 0$ $w = \frac{-1 \pm \sqrt{5}}{2}$ $\cos \frac{2\pi}{5} = \frac{\sqrt{5} - 1}{4}$ with reasons for choice	B1 B1 M1 A1 A1 E1	6	
	<b>Total</b>		<b>14</b>	
	<b>TOTAL</b>		<b>75</b>	